



**ROMMA**  
PURE CHEMISTRY



# Table Of Content

- ▶ SpS™ Super Purity Solvents
- ▶ UpS™ Ultra Purity Solvents
- ▶ High Purity Solvent Mixes
- ▶ Hi-Dry® Anhydrous Solvents
- ▶ Hi-Dry® KF Karl Fischer Reagents and Calibrants
- ▶ SpR™ Super Purity Reagents
- ▶ BiO™ BioPure Solvents
- ▶ PUROM™ High Purity Process Solvents
- ▶ BiO™ BioPure Solvents
- ▶ SpA™ Super Purity Acids and Reagents
- ▶ UpA™ Ultra Purity Acids and Reagents
- ▶ High Purity Aqueous Mixes
- ▶ PrimAg® Reference Materials
- ▶ PrimAg®-TvR Traceable Volumetric Reagents
- ▶ SA™ Select Assured™ Reagents for General Use
- ▶ Technical Grade Solvents and Acids

## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

**Acetic Acid glacial** (see Acetic Acid)

**2-Aminoethanol** (see Ethanolamine)

### Acetic Acid SpS

H014

500ml H014P  
1LT H014M  
2½LT H014L  
Dgr H:226-314  
P:280c-301+330+331-305+351+338-307+310



**(Acetic Acid glacial)**

CH<sub>3</sub>COOH MW 60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7]  
Assay >99.8% Water <0.1% Residue <0.0001%  
UV: 252nm >10%; 260nm >50%; 270nm >80%; 280nm >95%; 300-400nm >99%  
Application: Non-aqueous Titrations, Organic Trace Analysis (see also ROMIL Code H015 Acetic Acid SpA for inorganic trace analysis)

### Acetic Anhydride SpS

H027

500ml H027P  
1LT H027M  
2½LT H027L  
Dgr H:226-302+332-314-335  
P:210-280-301+330+331-305+351+338-317



**(CH<sub>3</sub>CO)<sub>2</sub>O** MW102.09FP-74°C BP 139°C d 1.08 CAS [108-24-7]

Assay >99% Residue <0.0001%  
Acetic Acid <1%  
Application: Molecular Biology

### Acetone SpS

H031

500ml H031P  
1LT H031M  
2½LT H031L  
Dgr H:225-319-336-EUH066  
P:210-233-305+351+338



**(Propanone)**

(CH<sub>3</sub>)<sub>2</sub>CO MW 58.08 BP 56.1°C d 0.79 CAS [67-64-1]  
Assay >99.9% Water <0.2% Residue <0.0001%  
UV: 329nm >10%; 335nm >50%; 340nm >80%; 345nm >95%; 350-400nm >99%  
Pesticide Residue Analysis passes test  
Application: HPLC, GC (eg, analysis of chlorohydrocarbons), UV, Environment Analysis (eg, pesticide residues)

### Acetone SpS

low in methanol

H032

2½LT H032L  
Dgr H:225-319-336-EUH066  
P:210-233-305+351+338



**(Propanone)**

(CH<sub>3</sub>)<sub>2</sub>CO MW 58.08 BP 56.1°C d 0.79 CAS [67-64-1]  
Assay >99.9% Water <0.2% Residue <0.0001%  
Methanol <10 ppm  
Application: GC, Applications requiring low methanol background

### Acetonitrile 200 SpS

far UV

H048

500ml H048P  
1LT H048M  
2½LT H048L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233



**(Methyl Cyanide)**

CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.01% Residue <0.0001%  
UV: 190nm >10%; 195nm >50%; 200nm >80%; 225nm >95%; 240-400nm >99%  
Pesticide Residue Analysis passes test  
Application: HPLC, Environment Analysis (eg, pesticide residues)

### Acetonitrile 190 SpS

far UV/gradient quality

H049

500ml H049P  
1LT H049M  
2½LT H049L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233



**(Methyl Cyanide)**

CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.005% Residue <0.0001%  
UV: 190nm >18%; 193nm >50%; 197nm >80%; 215nm >95%; 230-400nm >99%  
Acidity <0.0005 meq/g  
Alkalinity <0.00006 meq/g  
HPLC Gradient Use Test:  
205nm <0.005 AU  
254nm <0.002 AU  
Conforms to ACS liquid chromatography suitability.  
Pesticide Residue Analysis passes test  
IR Spectroscopy passes test  
Fluorescence Spectroscopy passes test  
Application: Gradient HPLC, GC, UV, IR, Environment Analysis (eg, pesticide residues), Molecular Biology (see also ROMIL Codes H051, H053, H055)  
Acetonitrile BiO very dry, Ion Chromatography using coulometric detection, Fluorescence Spectroscopy



## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

**2-Butanone** (see Methyl Ethyl Ketone)

**n-Butyl Chloride** (see 1-Chlorobutane)

**tert-Butyl Methyl Ether** (see Methyl tert-Butyl Ether)

500ml H083P  
1LT H083M  
2½LT H083L  
Dgr H:226-302-315-318-335-336  
P:210-280f-302+352-304+340-  
305+351+338-313



500ml H087P  
1LT H087M  
2½LT H087L  
Wng H:226-336-EUH066  
P:210



500ml H095P  
Dgr H:225-361fd-372-319-315  
P:210-233-280-302+352-  
305+351+338-403+235



500ml H104P  
1LT H104M  
2½LT H104L  
Wng H:226-332-315-411  
P:210-273-302+352-304+340



500ml H118P  
1LT H118M  
2½LT H118L  
Dgr H:225  
P:210



500ml H140P  
1LT H140M  
2½LT H140L  
Dgr H:351-361d-331-302-372-319-  
315  
P:261v-280f-304+340-305+351+338-  
308+313



### Butan-1-ol SpS

H083

**(n-Butanol, n-Butyl Alcohol)**

CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>OH MW 74.12 BP 117.7°C d 0.81 CAS [71-36-3]

Assay >99.8% Water <0.05% Residue <0.0001%

UV: 230nm >10%; 235nm >50%; 240nm >80%; 270nm >95%; 290-400nm >99%

### n-Butyl Acetate SpS

H087

CH<sub>3</sub>COO(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub> MW 116.16 BP 126.1°C d 0.88 CAS [123-86-4]

Assay >99.7% Water <0.05% Residue <0.0001%

UV: 257nm >10%; 260nm >50%; 275nm >80%; 310nm >95%; 360-400nm >99%

### Carbon Disulphide SpS

low in benzene

H095

CS<sub>2</sub> MW 76.13 BP 46.2°C d 1.26 CAS [75-15-0] Assay >99.9% Water <0.05% Residue <0.0005%

UV: 385nm >10%; 390nm >50%; 400nm >80%; 410nm >90%; 450nm >99%

Aromatic impurities (as benzene) <0.0030% (<30 ppm)

IR Spectroscopy passes test

Application: GC, IR, Environment Analysis (eg, determination of aromatics in air)

### Chlorobenzene SpS

H104

C<sub>6</sub>H<sub>5</sub>Cl MW 112.56 BP 131.7°C d 1.11 CAS [108-90-7]

Assay >99.9% Water <0.01% Residue <0.0001%

UV: 290nm >50%; 295nm >80%; 340nm >95%; 370-400nm >99%

Application: HPLC, UV Spectroscopy

### 1-Chlorobutane SpS

H118

**(n-Butyl Chloride)**

CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>Cl MW 92.57 BP 78.4°C d 0.88 CAS [109-69-3]

Assay >99.9% Water <0.005% Residue <0.0001%

UV: 235nm >10%; 240nm >50%; 250nm >80%; 255nm >95%; 290-400nm >99%

Application: HPLC, Molecular Biology (eg, protein sequencing)

### Chloroform SpS

stabilised with amylene

H140

**(Trichloromethane)**

CHCl<sub>3</sub> MW 119.38 BP 61.2°C d 1.48 CAS [67-66-3]

Assay >99.9%\* Water <0.005% Residue <0.0001%

UV: 250nm >10%; 255nm >50%; 260nm >80%; 270nm >95%; 280-400nm >99%

\*ex stabiliser

Pesticide Residue Analysis passes test

Stabiliser: Amylene ca. 25 ppm

Application: Gel Permeation Chromatography, Environment Analysis (eg, pesticide residues)



## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

### Chloroform SpS

stabilised with ethanol

H135

500ml H135P  
1LT H135M  
2½LT H135L  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280f-304+340-305+351+338-308+313



(Trichloromethane)

CHCl<sub>3</sub> MW 119.38 BP 61.2°C d 1.48CAS [67-66-3]  
Assay >99.9%\*Water <0.005% Residue <0.0001%  
UV: 250nm >10%; 255nm >50%; 260nm >80%; 270nm >95%; 280-400nm >99%  
\*ex stabiliser  
Pesticide Residue Analysis passes test  
IR Spectroscopy passes test  
Stabiliser: Ethanol ca. 1% w/w  
Stabiliser should only be removed immediately before use by adsorption onto activated alumina.  
Application: HPLC, UV, IR, Environment Analysis (eg, pesticide residues)

### Cyclohexane SpS

H156

500ml H156P  
1LT H156M  
2½LT H156L  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-302+352-331-403+235



C<sub>6</sub>H<sub>12</sub> MW 84.16 FP 6.5°C BP 80.7°C d 0.78 CAS [110-82-7]  
Assay >99.9% Water <0.005% Residue <0.0001%  
UV: 195nm >10%; 215nm >50%; 225nm >80%; 240nm >95%; 265-400nm >99%  
Pesticide Residue Analysis passes test  
Fluorescence Spectroscopy passes test  
Application: HPLC, UV, Fluorescence, Environment Analysis (eg, pesticide residues)

### Cyclohexanone SpS

H173

500ml H173P  
1LT H173M  
2½LT H173L  
Wng H:226-332  
P:210



C<sub>6</sub>H<sub>10</sub>OMW 98.15 FP-47°C BP 155°C d 0.94 CAS [108-94-1]  
Assay >99.8% Water <0.02% Residue <0.0001%  
UV: 340nm >10%; 345nm >50%; 350nm >80%; 365nm >95%; 390-400nm >99%  
Application:UV Spectroscopy

### 1,2-Dichlorobenzene SpS

H177

500ml H177P  
1LT H177M  
2½LT H177L  
Wng H:302-315-319-335-410  
P:273-302+352-305+351+338



C<sub>6</sub>H<sub>4</sub>Cl<sub>2</sub>MW 147.00 FP-17.0°C BP180.5°C d 1.31 CAS [95-50-1]  
Assay >99.8% Water <0.05% Residue <0.0005%  
UV: 300nm >10%; 310nm >50%; 330nm >80%; 375nm >95%; 390-400nm >99%  
Application: Gel Permeation Chromatography

### Dichloromethane SpS

stabilised with amylene

H202

500ml H202P  
1LT H202M  
2½LT H202L  
Wng H:351  
P:281-308+313



(Methylene Dichloride)

CH<sub>2</sub>Cl<sub>2</sub>MW 84.93 BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.01% Residue <0.0001%  
UV: 235nm >10%; 240nm >50%; 245nm >80%; 250nm >95%; 265-400nm >99%  
\*ex stabiliser  
IR Spectroscopy passes test  
Fluorescence Spectroscopy passes test  
Pesticide Residue Analysis passes test  
Stabiliser: Amylene ca. 25 ppm  
Application: HPLC, UV, IR, Fluorescence Analysis (eg, plasma cortisol), Environment Analysis (eg, pesticide residues), Molecular Biology (see also ROMILCode H203 Dichloromethane BiO), Gel Permeation Chromatography

### Diethyl Ether SpS

stabilised with BHT

H220

500ml H220P  
1LT H220M  
2½LT H220L  
Dgr H:224-302-336-EUH019-EUH066  
P:210-240-403+235



(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7]  
Assay >99.9%\* Water <0.02% Residue <0.0001%\*  
UV: 225nm >10%; 235nm >50%; 255nm >80%; 290nm >95%; 295-400nm >99%  
\*ex stabiliser  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 5 ppm  
Application: UVSpectroscopy, Techniques requiring a non-polar stabiliser

## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

500ml H218P 1LT H218M 2½LT H218L Dgr H:224-302-336-EUH019-EUH066 P:210-240-403+235	<h3>Diethyl Ether SpS</h3> <p>stabilised with copper <span style="float: right;">H218</span></p> <hr/> <p>(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7] □</p> <p>Assay &gt;99.9% Water &lt;0.02% Residue &lt;0.0001%</p> <p>UV: 220nm &gt;10%; 230nm &gt;50%; 250nm &gt;80%; 280nm &gt;95%; 315-400nm &gt;99%</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Stabiliser: Copper gauze</p> <p>Application: UV Spectroscopy, Separation applications requiring non-organic stabiliser</p>
500ml H219P 1LT H219M 2½LT H219L Dgr H:224-302-336-EUH019-EUH066 P:210-240-403+235	<h3>Diethyl Ether SpS</h3> <p>stabilised with ethanol <span style="float: right;">H219</span></p> <hr/> <p>(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7] □</p> <p>Assay &gt;99.9%* Water &lt;0.02% Residue &lt;0.0001%</p> <p>UV: 220nm &gt;10%; 230nm &gt;50%; 250nm &gt;80%; 270nm &gt;95%; 285-400nm &gt;99%</p> <p>*ex stabiliser</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Stabiliser: Ethanol ca. 1% v/v</p> <p>Application: HPLC, UV Spectroscopy, Applications requiring dry ether</p>
500ml H261P 1LT H261M 2½LT H261L Dgr H:225-360FD-332-EUH019 P:201-210-308+313-403+235	<h3>1,2-Dimethoxyethane SpS</h3> <p style="text-align: right;">H261</p> <hr/> <p>(Ethylene Glycol Dimethyl Ether) □</p> <p>CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> MW 90.12 FP -58°C BP 85°C d 0.87 CAS [110-71-4]</p> <p>Assay &gt;99.9% Water &lt;0.01% Residue &lt;0.0001%</p> <p>UV: 215nm &gt;10%; 240nm &gt;50%; 260nm &gt;80%; 280nm &gt;95%; 300-400nm &gt;99%</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Application: HPLC, UV Spectroscopy</p>
500ml H249P 1LT H249M 2½LT H249L Dgr H:227-360D-312+332-319 P:201-302+352-305+351+338-308+313	<h3>Dimethylacetamide SpS</h3> <p style="text-align: right;">H249</p> <hr/> <p>CH<sub>3</sub>CON(CH<sub>3</sub>)<sub>2</sub> MW 87.12 FP -20.0°C BP 166.1°C d 0.94 CAS [127-19-5]</p> <p>Assay &gt;99.7% Water &lt;0.05% Residue &lt;0.0001%</p> <p>UV: 275nm &gt;10%; 280nm &gt;50%; 300nm &gt;80%; 330nm &gt;95%; 350-400nm &gt;99%</p> <p>Application: Spectroscopy</p>
500ml H253P 1LT H253M 2½LT H253L Dgr H:360D-226-312+332-319 P:201-210-302+352-305+351+338-308+313	<h3>Dimethylformamide SpS</h3> <p style="text-align: right;">H253</p> <hr/> <p>HCON(CH<sub>3</sub>)<sub>2</sub> MW 73.09 BP 153.0°C d 0.95 CAS [68-12-2] □</p> <p>Assay &gt;99.9% Water &lt;0.03% Residue &lt;0.0001%</p> <p>UV: 270nm &gt;10%; 275nm &gt;50%; 285nm &gt;80%; 310nm &gt;95%; 325-400nm &gt;99%</p> <p>Application: HPLC, GC, Spectroscopy, Molecular Biology (see also ROMIL Code H254 Dimethylformamide BiO with 4Å molecular sieve), Gel Permeation Chromatography</p>
500ml H297P 1LT H297M 2½LT H297L Dgr H:225-350-319-335-EUH019-EUH066 P:210-281-305+351+338-308+313	<h3>1,4-Dioxan SpS</h3> <p style="text-align: right;">H297</p> <hr/> <p>C<sub>4</sub>H<sub>8</sub>O<sub>2</sub> MW 88.11 FP 11.8°C BP 101.3°C d 1.03 CAS [123-91-1] □</p> <p>Assay &gt;99.9% Water &lt;0.005% Residue &lt;0.0001%</p> <p>UV: 220nm &gt;10%; 240nm &gt;50%; 250nm &gt;80%; 280nm &gt;95%; 300-400nm &gt;99%</p> <p>Unstabilised</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Application: HPLC, GC, UV Spectroscopy, Liquid Scintillation, Applications requiring dry solvent</p>
500ml H236P 1LT H236M 2½LT H236L Dgr H:225-336-EUH019-EUH066 P:210-240-403+235	<h3>Di-iso-propyl Ether SpS</h3> <p>stabilised with BHT <span style="float: right;">H236</span></p> <hr/> <p>[(CH<sub>3</sub>)<sub>2</sub>CH]<sub>2</sub>O MW 102.18 BP 68.5°C d 0.73 CAS [108-20-3] □</p> <p>Assay &gt;99.5%* Water &lt;0.02% Residue &lt;0.0001%*</p> <p>UV: 245nm &gt;10%; 250nm &gt;50%; 260nm &gt;80%; 310nm &gt;95%; 330-400nm &gt;99%</p> <p>*ex stabiliser</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Stabiliser: Butylated hydroxytoluene (BHT) ca. 5 ppm</p> <p>Application: Spectroscopy</p>

## ROMIL-SpS™ Super Purity Solvents Specifications



high purity solvents for instrumental analysis

Ethyl Alcohol (see Ethanol)

Ethylene Glycol Dimethyl Ether (see 1,2-Dimethoxyethane)

### Ethanol absolute SpS

H314

500ml H314P (Ethyl Alcohol)  

1LT H314M C<sub>2</sub>H<sub>5</sub>OHW 46.07 BP 78.3°C d 0.79 CAS [64-17-5]

2½LT H314L Assay >99.8% Water <0.1% Residue <0.0001%

Dgr H:225 UV: 205nm >10%; 220nm >50%; 240nm >80%; 250nm >95%; 270-400nm >99%



P:210-233-240-403+235 IR Spectroscopy passes test

Fluorescence Spectroscopy passes test

Application: HPLC, GC, UV & IR & Fluorescence Spectroscopy (eg. uorimetric analysis of 17-ketosteroids)

### Ethanolamine SpS

H321

100ml H321S (2-Aminoethanol)  

500ml H321P CH<sub>2</sub>(OH)CH<sub>2</sub>NH<sub>2</sub> MW 61.08 FP 10.5°C BP 171°C d 1.02 CAS [141-43-5]

Dgr H:302+312+332-314 Assay >99.9% Water <0.2% Residue <0.0001%

P:280c-301+330+331-302+352-304+340-305+351+338-309+310 UV: 250nm >10%; 255nm >50%; 270nm >80%; 310nm >95%; 350-400nm >99%

Application: Molecular Biology, Liquid Scintillation

### Ethyl Acetate SpS

H346



500ml H346P CH<sub>3</sub>COOC<sub>2</sub>H<sub>5</sub> MW 88.11 BP 77.1°C d 0.90 CAS [141-78-6]

1LT H346M Assay >99.9% Water <0.005% Residue <0.0001%

2½LT H346L UV: 255nm >10%; 260nm >50%; 265nm >80%; 270nm >95%; 310-400nm >99%

Dgr H:225-319-336-EUHO66 Pesticide Residue Analysis passes test

P:210-233-240-305+351+338-403+235 Application: HPLC, GC, UV Spectroscopy, Environment Analysis (eg. pesticide residues), Molecular Biology

1,1,1,3,3,3-Hexa uoropropan-2-ol (see Hexa uoropropan-2-ol)

### Heptane fraction SpS

H368





500ml H368P C<sub>7</sub>H<sub>16</sub> BP 85-99°C d 0.69

1LT H368M Water <0.005% Residue <0.0001%

2½LT H368L UV: 195nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%


Dgr H:225-304-315-336-410 Comprises ca. 20-50% n-isomer, the remainder being predominantly other isomers of heptane.

P:210-273-301+310-331-302+352-304+340-403+235 Pesticide Residue Analysis passes test

    Application: HPLC, GC, UV Spectroscopy, Environment Analysis (eg. pesticide residues)

### n-Heptane 95% SpS

H367





500ml H367P CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW 100.21BP94-98°C d 0.68 CAS [142-82-5]  Water <0.005%

1LT H367M Residue <0.0001%

2½LT H367L UV: 195nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%

Dgr H:225-304-315-336-410 Assay (n-isomer) >95%

P:210-273-301+310-331-302+352-304+340-403+235 Assay (all isomers) >99.5%

    Pesticide Residue Analysis passes test

Fluorescence Spectroscopy passes test

Application: HPLC, GC, UV & Fluorescence Spectroscopy, Environment Analysis (eg.pesticide residues)

### n-Heptane 99% SpS

H366





500ml H366P CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>MW 100.21 BP98.4°C d 0.68 CAS [142-82-5]

1LT H366M Assay >99% Water <0.005% Residue <0.0001%

2½LT H366L UV: 195nm >10%; 210nm >50%; 220nm >80%; 245nm >95%; 290-400nm >99%



Dgr H:225-304-315-336-410 Application: GC, UV Spectroscopy, Molecular Biology

P:210-273-301+310-331-302+352-304+340-403+235

### Hexa uoropropan-2-ol SpS

H359

25ml H359V (1,1,1,3,3,3-Hexa uoropropan-2-ol)  

100ml H359S (CF<sub>3</sub>)<sub>2</sub>CHOH MW 168.04 FP -4.2°C BP 59.1°C d 1.62 CAS [920-66-1]

Dgr H:290-302+332-314 Assay >99.9% Water <0.05% Residue <0.0001%






P:280c-301+330+331-305+351+338-309+310 UV: 190nm >10%; 220nm >50%; 280nm >80%; 300nm >95%; 310-400nm >99%

Application: UV Spectroscopy, GC derivatisation reagent, Molecular Biology



## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

500ml H390P 1LT H390M 2½LT H390L Dgr H:225-304-361f-373-315-336-411 P:210-240-273-301+310-331-302+352-403+235	<h3>Hexane fraction SpS <span style="float: right;">H390</span></h3>
	<p>C6H14 BP 65-70°C d 0.66 CAS[73513-42-5] <span style="float: right;">□</span>            Water &lt;0.005% Residue &lt;0.0001%            UV: 190nm &gt;10%; 205nm &gt;50%; 220nm &gt;80%; 235nm &gt;95%; 255-400nm &gt;99%            Comprises ca. 50% n-isomer, the remainder being predominantly other isomers of hexane.            Pesticide Residue Analysis passes test            Application: HPLC, GC, UV Spectroscopy, Environment Analysis (eg, pesticide residues)</p>
500ml H388P 1LT H388M 2½LT H388L Dgr H:225-304-315-336-411 P:233-273-301+310-331-302+352-403+235	<h3>iso-Hexane 95% SpS <span style="float: right;">H388</span></h3>
	<p>C6H14 MW 86.18 BP55-63°Cd 0.65 CAS [107-83-5] <span style="float: right;">□</span> Water &lt;0.005% Residue &lt;0.0001%            UV: 195nm &gt;10%; 205nm &gt;50%; 220nm &gt;80%; 240nm &gt;95%; 260-400nm &gt;99%            n-Hexane &lt;5%            Pesticide Residue Analysis passes test            Fluorescence Spectroscopy passes test            Application: HPLC, GC, UV &amp; Fluorescence Spectroscopy, Environment Analysis (eg,pesticide residues)</p>
500ml H389P 1LT H389M 2½LT H389L Dgr H:225-304-361f-373-315-336-411 P:210-240-273-301+310-331-302+352-403+235	<h3>n-Hexane 95% SpS <span style="float: right;">H389</span></h3>
	<p>CH3(CH2)4CH3 MW86.18 BP67-70°C d 0.66 CAS [110-54-3] <span style="float: right;">□</span>            Water &lt;0.005% Residue &lt;0.0001%            UV: 190nm &gt;10%; 205nm &gt;50%; 220nm &gt;80%; 235nm &gt;95%; 255-400nm &gt;99%            Assay (n-isomer) &gt;95%            Assay (all isomers) &gt;99.5%            Pesticide Residue Analysis passes test            Fluorescence Spectroscopy passes test            Application: HPLC, GC, UV &amp; Fluorescence Spectroscopy, Environment Analysis (eg, pesticide residues)</p>
500ml H393P 1LT H393M 2½LT H393L Dgr H:225-304-361f-373-315-336-411 P:210-240-273-301+310-331-302+352-403+235	<h3>n-Hexane 99% SpS <span style="float: right;">H393</span></h3>
	<p>CH3(CH2)4CH3 MW 86.18 BP 68.7°C d 0.66 CAS [110-54-3] <span style="float: right;">□</span>            Assay &gt;99% Water &lt;0.005% Residue &lt;0.0001%            UV: 190nm &gt;10%; 205nm &gt;50%; 220nm &gt;80%; 235nm &gt;95%; 255-400nm &gt;99%            Pesticide Residue Analysis passes test            Application: HPLC, GC, UV Spectroscopy, Environment Analysis (eg, pesticide residues), Gel Permeation Chromatography</p>
<p><b>Methyl Alcohol (see Methanol)</b></p>	
<p><b>Methyl Cyanide (see Acetonitrile)</b></p>	
<p><b>4-Methyl-1,3-dioxolan-2-one (see Propylene Carbonate)</b></p>	
<p><b>Methylene Chloride (see Dichloromethane)</b></p>	
<p><b>Methylene Dichloride (see Dichloromethane) 4-</b></p>	
<p><b>Methylpentan-2-one (see Methyl iso-Butyl Ketone)</b></p>	
500ml H409P 1LT H409M 2½LT H409L Dgr H:225-301+311+331-370 P:210-280f-302+352-309+310-403+235	<h3>Methanol 215 SpS <span style="float: right;">H409</span></h3>
	<p>(Methyl Alcohol) <span style="float: right;">□</span>            CH3OHMW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]            Assay &gt;99.9% Water &lt;0.02% Residue &lt;0.0001%            UV: 215nm &gt;10%; 225nm &gt;50%; 235nm &gt;80%; 255nm &gt;95%; 280-400nm &gt;99%            Application: HPLC, UV, Liquid Scintillation</p>


## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

### Methanol 205 SpS


gradient quality

H410

<p>500ml H410P (Methyl Alcohol)</p> <p>1LT H410M</p> <p>2½LT H410L</p> <p>Dgr H:225-301+311+331-370</p> <p>P:210-280F-302+352-309+310-403+235</p> 	<p>CH<sub>3</sub>OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]</p> <p>Assay &gt;99.9% Water &lt;0.02% Residue &lt;0.0001%</p> <p>UV: 205nm &gt;10%; 210nm &gt;50%; 225nm &gt;80%; 240nm &gt;95%; 265-400nm &gt;99%</p> <p>Acidity &lt;0.0003 meq/g</p> <p>Alkalinity &lt;0.0002 meq/g</p> <p>HPLC Gradient Use Test:</p> <p>230nm &lt;0.005 AU</p> <p>254nm &lt;0.005 AU</p> <p>Conforms to ACS liquid chromatography suitability.</p> <p>Pesticide Residue Analysis passes test</p> <p>Fluorescence Spectroscopy passes test</p> <p>Application: Gradient HPLC, GC, UV, Environment Analysis (eg, pesticide residues), Applications requiring dry solvent, Non-aqueous Titrations, Liquid Scintillation, Molecular Biology, Fluorescence Spectroscopy</p>
---	---


### Methyl tert-Butyl Ether SpS

H447

<p>500ml H447P</p> <p>1LT H447M</p> <p>2½LT H447L</p> <p>Dgr H:225-315</p> <p>P:210-233-302+352-403+235</p> 	<p>(tert-Butyl Methyl Ether)</p> <p>CH<sub>3</sub>OC(CH<sub>3</sub>)<sub>3</sub> MW 88.15 BP 55.4°Cd 0.74CAS [1634-04-4]</p> <p>Assay&gt;99.7% Water &lt;0.02% Residue &lt;0.0001%</p> <p>UV: 220nm &gt;10%; 235nm &gt;50%; 255nm &gt;80%; 270nm &gt;90%; 300-400nm &gt;99%</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Application: HPLC, GC</p>
---	--


### Methyl iso-Butyl Ketone SpS

H446

<p>500ml H446P</p> <p>1LT H446M</p> <p>2½LT H446L</p> <p>Dgr H:225-332-319-336-351-EUH066</p> <p>P:210-305+351+338-304+340</p> 	<p>(4-Methylpentan-2-one)</p> <p>(CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>COCH<sub>3</sub> MW 100.16 BP 117.4°C d 0.80 CAS [108-10-1]</p> <p>Assay &gt;99.8% Water &lt;0.01% Residue &lt;0.0001%</p> <p>UV: 335nm &gt;10%; 340nm &gt;50%; 360nm &gt;80%; 375nm &gt;95%; 390-400nm &gt;99%</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Application: Organic Analysis and Chromatography</p>
--	--


### Methylcyclohexane SpS

H465

<p>500ml H465P</p> <p>1LT H465M</p> <p>Dgr H:225-304-315-336-411</p> <p>P:273-301+310-331-302+352-403+235</p> 	<p>C<sub>7</sub>H<sub>14</sub> MW 98.19 BP 101°Cd 0.77CAS [108-87-2]</p> <p>Assay &gt;99.9% Water &lt;0.005% Residue &lt;0.0001%</p> <p>UV: 200nm &gt;10%; 220nm &gt;50%; 230nm &gt;80%; 250nm &gt;95%; 280-400nm &gt;99%</p> <p>Fluorescence Spectroscopy passes test</p> <p>Application: UV Spectroscopy, Fluorescence Spectroscopy</p>
---	---


### Methyl Ethyl Ketone SpS

H493

<p>500ml H493P</p> <p>1LT H493M</p> <p>2½LT H493L</p> <p>Dgr H:225-319-336-EUH066</p> <p>P:210-305+351+338-403+233</p> 	<p>(2-Butanone)</p> <p>CH<sub>3</sub>CH<sub>2</sub>COCH<sub>3</sub> MW 72.11 BP 79.6°C d 0.80 CAS [78-93-3]</p> <p>Assay &gt;99.8% Water &lt;0.02% Residue &lt;0.0001%</p> <p>UV: 330nm &gt;10%; 335nm &gt;50%; 340nm &gt;80%; 345nm &gt;95%; 350-400nm &gt;99%</p> <p>Application: HPLC, UV Spectroscopy, Environment Analysis</p>
--	---


### N-Methyl-2-pyrrolidone SpS

H565

<p>500ml H565P</p> <p>1LT H565M</p> <p>2½LT H565L</p> <p>Dgr H:360D-315-319-335</p> <p>P:201-302+352-305+351+338-308+313</p> 	<p>CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONCH<sub>3</sub> MW 99.13 BP 202.0°C d 1.03 CAS [872-50-4]</p> <p>Assay &gt;99.5% Water &lt;0.05%</p> <p>UV: 295nm &gt;10%; 300nm &gt;50%; 320nm &gt;80%; 360nm &gt;95%; 395-400nm &gt;99%</p> <p>Application: Molecular Biology (eg, DNA synthesis), Versatile and powerful solvent properties</p>
--	---

### 2-Methyltetrahydrofuran SpS

H536

<p>500ml H536P</p> <p>1LT H536M</p> <p>2½LT H536L</p> <p>Dgr H:225-319-335-EUH019</p> <p>P:210-233-240-305+351+338-403+235</p> 	<p>CH<sub>3</sub>C<sub>4</sub>H<sub>7</sub>O MW 86.13 BP 80°C d 0.86 CAS [96-47-9]</p> <p>Assay &gt;99.8% Water &lt;0.01% Residue &lt;0.0001%</p> <p>UV: 245nm &gt;10%; 250nm &gt;50%; 270nm &gt;80%; 295nm &gt;95%; 360-400nm &gt;99%</p> <p>Unstabilised</p> <p>Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm)</p> <p>Application: Applications requiring dry solvent</p>
--	---

## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

### 2-Methyltetrahydrofuran SpS

stabilised with BHT

H537

500ml H537P  
1LT H537M  
2½LT H537L  
Dgr H:225-319-335-EUH019  
P:210-233-240-305+351+338-403+235



CH<sub>3</sub>C<sub>4</sub>H<sub>7</sub>O MW 86.13 BP 80°C d 0.86 CAS [96-47-9] □  
Assay >99.8%\* Water <0.01% Residue <0.0001%\*  
\*ex stabiliser  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm  
Application: Applications requiring dry solvent

### n-Nonane 95% SpS

H 5 6 8

500ml H568P  
1LT H568M  
2½LT H568L  
Dgr H:226-304  
P:210-301+310-331



CH<sub>3</sub>(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub> MW128.26BP146-150°C d 0.72 CAS [111-84-2] □  
Water <0.005% Residue <0.0001%  
UV: 220nm >10%; 240nm >50%; 290nm >80%; 310nm >95%; 330-400nm >99%  
Application: UV Spectroscopy

[iso-Octane \(see 2,2,4-Trimethylpentane\)](#)

[Perchloroethylene \(see Tetrachloroethylene\)](#)

[Petroleum Distillate \(see Petroleum Ether\)](#)

[n-Propanol \(see Propan-1-ol\)](#)

[iso-Propanol \(see Propan-2-ol\)](#)

[Propanone \(see Acetone\)](#)

[n-Propyl Alcohol \(see Propan-1-ol\)](#)

[iso-Propyl Alcohol \(see Propan-2-ol\)](#)

### n-Pentane 95% SpS

H571

500ml H571P  
1LT H571M  
2½LT H571L  
Dgr H:225-304-336-411-EUH066  
P:273-301+310-331-403+235



CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub> MW 72.15 BP35.5-37°C d 0.63 CAS [109-66-0] □  
Water <0.005% Residue <0.0001%  
UV: 190nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%  
Assay (n-isomer) >95%  
Assay (all isomers) >99.5%  
Haloform impurities <0.0001% (<1 ppm)  
Pesticide Residue Analysis passes test  
Application: HPLC, GC, Environment Analysis (eg pesticide residues), Gel Permeation Chromatography

### Petroleum Ether 30-40°C SpS

H600

500ml H600P  
1LT H600M  
2½LT H600L  
Dgr H:225-304-336-411-EUH066  
P:210-233-243-273-280-301+310-303+361+353-405-501



**(Petroleum Distillate, Petroleum Spirit)** □  
BP 30-40°C d 0.64 CAS [109-66-0]  
Water <0.005% Residue <0.0001%  
UV: 190nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%

### Petroleum Ether 40-60°C SpS

H601

500ml H601P  
1LT H601M  
2½LT H601L  
Dgr H:225-304-336-411-EUH066  
P:210-233-243-273-280-301+310-303+361+353-304+340-331-403+235



**(Petroleum Distillate, Petroleum Spirit)** □  
BP 40-60°C d 0.64 CAS [8032-32-4]  
Water <0.005% Residue <0.0001%  
UV: 190nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%  
Pesticide Residue Analysis passes test  
Application: Environment Analysis (eg pesticide residues)



## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

500ml H602P  
1LT H602M  
2½LT H602L  
Dgr H:225-304-315-336-411  
P:210-243-273-280-301+310-331-403+235



### Petroleum Ether 60-80°C SpS

H602

**(Petroleum Distillate, Petroleum Spirit)**

BP 60-80°C d 0.67  
Water <0.005% Residue <0.0001%  
UV: 190nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%  
Pesticide Residue Analysis passes test  
Application: Environment Analysis (eg pesticide residues)

500ml H603P  
1LT H603M  
2½LT H603L  
Dgr H:225-304-315-336-411  
P:210-273-280-260v-301+310-331-403+235



### Petroleum Ether 80-100°C SpS

H603

**(Petroleum Distillate, Petroleum Spirit)**

BP 80-100°C d 0.69CAS [64742-49-0]  
Water <0.005% Residue <0.0001%  
UV: 200nm >10%; 215nm >50%; 230nm >80%; 240nm >95%; 255-400nm >99%

500ml H629P  
1LT H629M  
2½LT H629L



### Propan-1,2-diol SpS

H629

**(1,2-Propylene Glycol)**

CH<sub>3</sub>CH(OH)CH<sub>2</sub>OH MW 76.10 BP 187.6°C CAS [57-55-6]  
Assay >99.8% Water <0.005% Residue <0.0001%  
UV: 210nm >10%; 215nm >50%; 230nm >80%; 245nm >95%; 260-400nm >99%  
Application: UV Spectroscopy

500ml H624P  
1LT H624M  
2½LT H624L  
Dgr H:225-318-336  
P:210-233-280f-305+351+338-313



### Propan-1-ol SpS

H624

**(n-Propanol, n-Propyl Alcohol)**

CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>OH MW 60.10 BP 97.2°C d 0.80 CAS [71-23-8]  
Assay >99.9% Water <0.05% Residue <0.0001%  
UV: 210nm >10%; 225nm >50%; 235nm >80%; 250nm >95%; 280-400nm >99%  
Application: HPLC, GC, UV Spectroscopy

500ml H625P  
1LT H625M  
2½LT H625L  
Dgr H:225-319-336  
P:210-233-305+351+338



### Propan-2-ol SpS

H625

**(iso-Propanol, iso-Propyl Alcohol)**

(CH<sub>3</sub>)<sub>2</sub>CHOH MW 60.10BP 82.2°C d 0.78 CAS [67-63-0]  
Assay >99.9% Water <0.02% Residue <0.0001%  
UV: 205nm >10%; 210nm >50%; 225nm >80%; 240nm >95%; 255-400nm >99%  
Pesticide Residue Analysis passes test  
Fluorescence Spectroscopy passes test  
Application: HPLC, GC, UV & Fluorescence Spectroscopy, Environment Analysis (eg, pesticide residues)

500ml H645P  
1LT H645M  
Wng H:319  
P:305+351+338



### Propylene Carbonate SpS

H645

**(4-Methyl-1,3-dioxolan-2-one)**

CH<sub>3</sub>CHOCOOCH<sub>2</sub> MW102.09 BP 241.7°C d 1.20 CAS [108-32-7]  
Assay >99.8% Water <0.01% Residue <0.0001%  
UV: 255nm >10%; 290nm >50%; 315nm >80%; 350nm >95%; 360-400nm >99%  
Application: GC

500ml H650P  
1LT H650M  
2½LT H650L  
Dgr H:225-302+312+332-315-319  
P:210-302+352-304+340-305+351+338-403+235



### Pyridine SpS

H650

C<sub>5</sub>H<sub>5</sub>N MW 79.10 BP 115.3°C d 0.98 CAS [110-86-1]  
Assay >99.8% Water <0.02% Residue <0.0001%  
UV: 330nm >10%; 335nm >50%; 340nm >80%; 350nm >95%; 370-400nm >99%  
IR Spectroscopy passes test  
Application: UV and IR Spectroscopy

500ml H702P  
1LT H702M  
2½LT H702L  
Wng H:315-317-319-336-351-411  
P:273-281-302+352-305+351+338-308+313



### Tetrachloroethylene SpS

H702

**(Perchloroethylene)**

CCl<sub>2</sub>CCl<sub>2</sub> MW 165.83 BP 121.1°C d 1.62 CAS [127-18-4]  
Assay >99.9% Water <0.005% Residue <0.0001%  
UV: 290nm >10%; 295nm >50%; 300nm >80%; 380nm >95%; 395-400nm >99%  
Unstabilised  
IR Spectroscopy passes test  
Application: UV & IR Spectroscopy

## ROMIL-SpS™ Super Purity Solvents Specifications


high purity solvents for instrumental analysis

500ml H718P  
1LT H718M  
2½LT H718L  
Dgr H:225-319-335-351-EUHO19  
P:210-240-305+351+338-308+313-403+233



### Tetrahydrofuran SpS

H718

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66.0°C d 0.89 CAS [109-99-9]   
Assay >99.9% Water <0.005% Residue <0.0001%  
UV: 215nm >10%; 235nm >50%; 255nm >80%; 275nm >95%; 295-400nm >99%  
Unstabilised  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Application: HPLC, Gel Permeation Chromatography, UV Spectroscopy, Applications requiring drysolvent, Molecular Biology


500ml H719P  
1LT H719M  
2½LT H719L  
Dgr H:225-319-335-351-EUHO19  
P:210-240-305+351+338-308+313-403+233



### Tetrahydrofuran SpS

stabilised with BHT

H719


CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66.0°C d 0.89 CAS [109-99-9]  Assay >99.9%\*  
Water <0.005% Residue <0.0001%\*  
UV: 290nm >10%; 295nm >50%; 300nm >80%; 305nm >95%; 310-400nm >99%  
\*ex stabiliser  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm  
Application: Applications requiring dry solvent, Analysis of vinyl chloride in PVC, Gel Permeation Chromatography

500ml H771P  
1LT H771M  
2½LT H771L  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235



### Toluene SpS

H771


C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub> MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]   
Assay >99.9% Water <0.01% Residue <0.0001%  
UV: 285nm >10%; 290nm >50%; 300nm >80%; 320nm >95%; 350-400nm >99%  
Pesticide Residue Analysis passes test  
Application: HPLC, GC, Environment Analysis (eg, pesticide residues), Gel Permeation Chromatography, Liquid Scintillation

500ml H742P  
1LT H742M  
2½LT H742L  
Dgr H:350-315-319-336-341-412  
P:201-273-302+352-305-351-338-308+313



### Trichloroethylene SpS

H742


CCl<sub>2</sub>CHCl MW 131.39 FP -87°C BP 86.7°C d 1.46 CAS [79-01-6]   
Assay >99.8%\* Water <0.01% Residue <0.0001%  
UV: 275nm >10%; 280nm >50%; 315nm >80%; 375nm >95%; 400nm >99%  
\*ex stabiliser  
Stabiliser: Maxistab ca. 700 ppm  
Application: UV Spectroscopy  
Maxistab is a trademark of The Dow Chemical Company

100ml H860S  
500ml H860P  
Dgr H:226-301+331-312-315-318-373  
P:210-280f-302+352-304+340-305+351+338-309+310



### 2,2,2-Tri uoroethanol SpS

H860


CF<sub>3</sub>CH<sub>2</sub>OH MW100.04BP 74.1°Cd 1.39 CAS [75-89-8]   
Assay >99.9% Water <0.1% Residue <0.0001%  
UV: 190nm >40%; 195nm >70%; 200nm >80%; 230nm >95%; 265-400nm >99%  
Application: UV Spectroscopy (very low cut o )

500ml H901P  
1LT H901M  
2½LT H901L  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-331-302+352-304+340-403+235



### 2,2,4-Trimethylpentane SpS


H901

(*iso*-Octane)   
(CH<sub>3</sub>)<sub>3</sub>CCCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1]  
Assay >99.75% Water <0.005% Residue <0.0001%  
UV: 205nm >10%; 215nm >50%; 225nm >80%; 240nm >95%; 270-400nm >99%  
IR Spectroscopy passes test  
Pesticide Residue Analysis passes test  
Application: HPLC, GC, UV & IR Spectroscopy, Environment Analysis (eg, pesticide residues)

500ml H950P  
1LT H950M  
2½LT H950L

### Water SpS

H 9 5 0

H<sub>2</sub>O MW 18.02 FP 0.0°C BP 100.0°C CAS [7732-18-5]   
Residue <0.0001%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
TOC (at time of manufacture) <30 ppb  
HPLC Gradient Use Test:  
205nm <0.005 AU  
254nm <0.002 AU  
Conforms to ACS liquid chromatography suitability.  
Filtered to 0.2 micron  
Application: Gradient HPLC, Environment Analysis of Trace Organics (see also ROMIL Code H951 Water SpA for inorganic trace analysis), Molecular Biology

## ROMIL-SpS™ Super Purity Solvents Specifications

high purity solvents for instrumental analysis

### m-Xylene SpS

H965

500ml H965P C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW 106.17 FP -47.8°C BP 139.1°C d 0.86 CAS [108-38-3] □  
 Wng H:226-312+332-315 Assay >99.0% Water <0.01% Residue <0.0001%  
 P:210-302+352-304+340 UV: 295nm >10%; 305nm >50%; 320nm >80%; 350nm >95%; 370-400nm >99%



### o-Xylene SpS

H958

500ml H958P C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW 106.17 FP -25°C BP 144.4°C d 0.88 CAS [95-47-6] □  
 Wng H:226-312+332-315 Assay >99.0% Water <0.01% Residue <0.0001%  
 P:210-302+352 UV: 295nm >10%; 305nm >50%; 320nm >80%; 350nm >95%; 370-400nm >99%



### p-Xylene SpS

H973

500ml H973P C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW 106.17 FP 13.2°C BP 138.3°C d 0.86 CAS [106-42-3] □  
 Wng H:226-312+332-315 Assay >99.0% Water <0.01% Residue <0.0001%  
 P:210-302+352 UV: 295nm >10%; 305nm >50%; 320nm >80%; 350nm >95%; 370-400nm >99%



### Xylene mixed isomers SpS

H982

500ml H982P C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW 106.17 BP 138-142°C d 0.86 CAS [1330-20-7] □  
 1LT H982M Water <0.01% Residue <0.0001%  
 2½LT H982L UV: 295nm >10%; 305nm >50%; 320nm >80%; 350nm >95%; 370-400nm >99%  
 Wng H:226-312+332-315 Comprises 3 isomers and ethylbenzene  
 P:210-302+352-304+340 Assay (total C<sub>8</sub>H<sub>10</sub> isomers) >98.5%  
 Ethylbenzene typically <3%  
 Toluene typically <0.5%  
 Methyl ethylbenzene typically <0.5%  
 Application: Liquid Scintillation





## ROMIL-UpS™ Ultra Purity Solvents Specifications

for critical analytical applications

### Acetone UpS

ultra gc

H033

2½LT H035L  
Dgr H:225-319-336-EUH066  
P:210-233-305+351+338



(Propanone)

(CH<sub>3</sub>)<sub>2</sub>CO MW 58.08 BP 56.1°C d 0.79 CAS [67-64-1]  
Assay >99.95% Water <0.2% Residue <0.0001%  
Suitability for GC-FID passes test  
Suitability for GC-ECD passes test  
Suitability for GC-MS passes test  
Application:GC critical lowresidue applications

### Acetonitrile UpS

ultra gc

H054

2½LT H054L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233



(Methyl Cyanide)

CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.02% Residue <0.0001%  
Suitability for GC-FID passes test  
Suitability for GC-ECD passes test  
Suitability for GC-MS passes test  
Application:GCcritical lowresidue applications

### Acetonitrile UpS

ultra lc

H050

1LT H050M  
2½LT H050L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233



(Methyl Cyanide)

CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.005% Residue <0.0001%  
UV: 190nm >18%; 193nm >50%; 197nm >80%; 215nm >95%; 230-400nm >99%  
Acidity <0.0005 meq/g  
Alkalinity <0.00006 meq/g  
Gradient Use Test: 205nm <0.002 AU; 254nm <0.0005 AU  
Baseline drift <0.02 AU @ 205nm  
Fluorescence (as quinine): 254nm <1 ppb; 365nm <1 ppb  
Suitability for LC-MS passes test  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Application: HPLC criticalgradient applications, LC-MS, UHPLC

### Acetonitrile UpS

ultra pfas

H052

2½LT H052L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233



(Methyl Cyanide)

CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.005% Residue <0.0001%  
UV: 190 nm >18%; 193nm >50%; 197 nm > 80%; 215nm >95%; 230-400nm >99%  
Acidity <0.0005 meq/g  
Alkalinity <0.00006 meq/g  
Gradient Use Test: 205nm <0.002 AU; 254nm <0.0005 AU  
Baseline drift <0.02 AU @ 205nm  
Fluorescence (as quinine): 254nm <1 ppb; 365nm <1 ppb  
Suitability for LC-MS passes test  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Suitability for PFAS analysis passes test  
Application: Ultra low background solvent for LC-MS analysis of poly- and per-  
uoroalkyl substances

### Dichloromethane UpS

stabilised with amylene ultra lc

H204

2½LT H204L  
Wng H:351  
P:281-308+313



(Methylene Dichloride)

CH<sub>2</sub>Cl<sub>2</sub>MW 84.93 BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.01% Residue <0.0001%  
UV: 235nm >10%; 240nm >50%; 245nm >80%; 250nm >95%; 265-400nm >99%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm  
Gradient Use Test: 260nm <0.0005 AU  
Application: HPLC critical gradient applications, UHPLC

### Dichloromethane UpS

stabilised with cyclohexene/amylen ultra gc

H205

2½LT H205L  
Wng H:351  
P:281-308+313






















(Methylene Dichloride)

CH<sub>2</sub>Cl<sub>2</sub>MW84.93BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.01% Residue <0.0001%  
\*ex stabiliser  
Stabiliser: Cyclohexene/Amylene ca. 50/25 ppm  
Suitability for GC-FID: passes test  
Suitability for GC-ECD: passes test  
Suitability for GC-MS: passes test  
Application: GC critical low residue applications

## ROMIL-UpS™ Ultra Purity Solvents Specifications

for critical analytical applications

<p>2½LT H317L Dgr H:225 P:210-233-240-403+235</p> 	<p><b>Ethyl Alcohol (see Ethanol)</b></p> <p><b>Ethanol absolute UpS</b> <b>ultra lc</b> <span style="float: right;">H317</span></p> <p><b>(Ethyl Alcohol)</b></p> <p>C<sub>2</sub>H<sub>5</sub>OH MW 46.07 BP 78.3°C d 0.79 CAS [64-17-5] Assay &gt;99.8% Water &lt;0.1% Residue &lt;0.0001% UV: 205nm &gt;10%; 220nm &gt;50%; 240nm &gt;80%; 250nm &gt;95%; 270-400nm &gt;99% Gradient Use Test: &lt;0.005 AU @ 260nm Fluorescence (as quinine): 254nm &lt;2 ppb Application: HPLC critical gradient applications, UHPLC</p>
<p>2½LT H347L Dgr H:225-319-336-EUH066 P:210-233-240-305+351+338-403+235</p>  	<p><b>Ethyl Acetate UpS</b> <b>ultra lc</b> <span style="float: right;">H347</span></p> <p>CH<sub>3</sub>COOC<sub>2</sub>H<sub>5</sub> MW 88.11 BP 77.1°C d 0.90 CAS [141-78-6] Assay &gt;99.9% Water &lt;0.005% Residue &lt;0.0001% UV: 255nm &gt;10%; 260nm &gt;50%; 265nm &gt;80%; 270nm &gt;95%; 310-400nm &gt;99% Gradient Use Test: &lt;0.0005 AU @ 290nm Application: HPLC critical gradient applications, UHPLC</p>
<p>2½LT H364L Dgr H:225-304-315-336-410 P:210-273-301+310-331-302+352-304+340-403+235</p>    	<p><b>n-Heptane 99% UpS</b> <b>ultra gc</b> <span style="float: right;">H364</span></p> <p>CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW 100.21 BP 98.4°C d 0.68 CAS [142-82-5] Assay &gt;99% Water &lt;0.01% Residue &lt;0.0001% Suitability for GC-FID passes test Suitability for GC-ECD passes test Suitability for GC-MS passes test Application: GC critical low residue applications</p>
<p>1LT H363M 2½LT H363L Dgr H:225-304-315-336-410 P:210-273-301+310-331-302+352-304+340-403+235</p>    	<p><b>n-Heptane 99% UpS</b> <b>ultra lc</b> <span style="float: right;">H363</span></p> <p>CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW 100.21 BP 98.4°C d 0.68 CAS [142-82-5] Assay &gt;99% Water &lt;0.005% Residue &lt;0.0001% UV: 195nm &gt;10%; 210nm &gt;50%; 220nm &gt;80%; 245nm &gt;95%; 290-400nm &gt;99% Acidity &lt;0.0002 meq/g Alkalinity &lt;0.0002 meq/g Trace ionic impurities: Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn &lt;10 ppb each Al, Ca &lt;25 ppb each Na &lt;50 ppb Application: HPLC critical applications, LC-MS, UHPLC</p>
<p>2½LT H391L Dgr H:225-304-361f-373-315-336-411 P:210-240-273-301+310-331-302+352-403+235</p>    	<p><b>n-Hexane 95% UpS</b> <b>ultra lc</b> <span style="float: right;">H391</span></p> <p>CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> MW 86.18 BP 67-70°C d 0.66 CAS [110-54-3] Water &lt;0.005% Residue &lt;0.0001% UV: 190nm &gt;10%; 205nm &gt;50%; 220nm &gt;80%; 235nm &gt;95%; 255-400nm &gt;99% Assay (n-isomer) &gt;95% Assay (all isomers) &gt;99.5% Gradient Use Test: &lt;0.0005 AU @ 260nm Application: HPLC critical gradient applications, UHPLC</p>
<p>2½LT H394L Dgr H:225-304-361f-373-315-336-411 P:210-240-273-301+310-331-302+352-403+235</p>    	<p><b>n-Hexane 99% UpS</b> <b>ultra gc</b> <span style="float: right;">H394</span></p> <p>CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> MW 86.18 BP 68.7°C d 0.66 CAS [110-54-3] Assay &gt;99% Water &lt;0.01% Residue &lt;0.0001% Suitability for GC-FID passes test Suitability for GC-ECD passes test Suitability for GC-MS passes test Application: GC critical low residue applications</p>

## ROMIL-UpS™ Ultra Purity Solvents Specifications

for critical analytical applications

### n-Hexane 99% UpS

ultra lc

H395

1LT H395M  
2½LT H395L  
Dgr H:225-304-361f-373-315-336-411  
P:210-240-273-301+310-331-302+352-403+235



CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> MW 86.18 BP 68.7°C d 0.66 CAS [110-54-3]  
Assay >99% Water <0.005% Residue <0.0001%  
UV: 190nm >10%; 205nm >50%; 220nm >80%; 235nm >95%; 255-400nm >99%  
Acidity <0.0002 meq/g  
Alkalinity <0.0002 meq/g  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Application: HPLC critical applications, LC-MS, UHPLC

**Methyl Alcohol** (see [Methanol](#))

**Methyl Cyanide** (see [Acetonitrile](#))

**Methylene Dichloride** (see [Dichloromethane](#))

### Methanol UpS

ultra gc

H415

2½LT H415L  
Dgr H:225-301+311+331-370  
P:210-280f-302+352-309+310-403+235



(Methyl Alcohol)  
CH<sub>3</sub>OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
Assay >99.9% Water <0.02% Residue <0.0001%  
Suitability for GC-FID passes test  
Suitability for GC-ECD passes test  
Suitability for GC-MS passes test  
Application: GC critical low residue applications

### Methanol UpS

ultra lc

H411

1LT H411M  
2½LT H411L  
Dgr H:225-301+311+331-370  
P:210-280f-302+352-309+310-403+235



(Methyl Alcohol)  
CH<sub>3</sub>OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
Assay >99.9% Water <0.02% Residue <0.0001%  
UV: 205nm >10%; 210nm >50%; 225nm >80%; 240nm >95%; 265-400nm >99%  
Acidity <0.0003 meq/g  
Alkalinity <0.0002 meq/g  
Gradient Use Test: 230nm <0.002 AU; 254nm <0.002 AU  
Baseline drift <0.02 @ 230nm  
Fluorescence (as quinine): 254nm <1 ppb; 365nm <1 ppb  
Suitability for LC-MS passes test  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Application: HPLC critical gradient applications, LC-MS, UHPLC

### Methanol UpS

ultra pfas

H414

2½LT H414L  
Dgr H:225-301+311+331-370  
P:210-280f-302+352-309+310-403+235



(Methyl Alcohol)  
CH<sub>3</sub>OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
Assay >99.9% Water <0.02% Residue <0.0001%  
UV: 205nm >10%; 210nm >50%; 225nm >80%; 240nm >95%; 265-400nm >99%  
Acidity <0.0003 meq/g  
Alkalinity <0.0002 meq/g  
Gradient Use Test: 230nm <0.002 AU; 254nm <0.002 AU  
Baseline drift <0.02 @ 230 nm  
Fluorescence (as quinine): 254nm <1 ppb; 365nm <1 ppb  
Suitability for LC-MS passes test  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Suitability for PFAS analysis passes test  
Application: Ultra low background solvent for LC-MS analysis of poly- and perfluoroalkyl substances

**iso-Octane** (see [2,2,4-Trimethylpentane](#))

**iso-Propanol** (see [Propan-2-ol](#))







**Propanone** (see [Acetone](#))

**iso-Propyl Alcohol** (see [Propan-2-ol](#))



## ROMIL-UpS™ Ultra Purity Solvents Specifications

for critical analytical applications

<p>2½LT H574L Dgr H:225-304-336-411-EUH066 P:273-301+310-331-403+235</p> 	<p><b>n-Pentane 95% UpS</b> <b>ultra gc</b> <span style="float: right;">H574</span></p> <p><chem>CH3(CH2)3CH3</chem> MW 72.15 BP 36.0°C d 0.63 CAS [109-66-0] Water &lt;0.005% Residue &lt;0.0001% Assay (n-isomer) &gt;95% Assay (all isomers) &gt;99.5% Suitability for GC-FID passes test Suitability for GC-ECD passes test Suitability for GC-MS passes test Application: GC critical low residue applications</p>
<p>2½LT H573L Dgr H:225-304-336-411-EUH066 P:273-301+310-331-403+235</p> 	<p><b>n-Pentane 99% UpS</b> <b>ultra gc</b> <span style="float: right;">H573</span></p> <p><chem>CH3(CH2)3CH3</chem> MW 72.15 BP 36.0°C d 0.63 CAS [109-66-0] Assay &gt;99% Water &lt;0.01% Residue &lt;0.0001% Suitability for GC-FID passes test Suitability for GC-ECD passes test Suitability for GC-MS passes test Application: GC critical low residue applications</p>
<p>1LT H626M 2½LT H626L Dgr H:225-319-336 P:210-233-305+351+338</p> 	<p><b>Propan-2-ol UpS</b> <b>ultra lc</b> <span style="float: right;">H626</span></p> <p><b>(iso-Propanol, iso-Propyl Alcohol)</b> <chem>(CH3)2CHOH</chem> MW 60.10 BP 82.2°C d 0.78 CAS [67-63-0] Assay &gt;99.9% Water &lt;0.02% Residue &lt;0.0001% UV: 205nm &gt;10%; 210nm &gt;50%; 225nm &gt;80%; 240nm &gt;95%; 255-400nm &gt;99% Acidity &lt;0.0002 meq/g Alkalinity &lt;0.0002 meq/g Gradient Use Test: 254nm &lt;0.005 AU Fluorescence (as quinine): 254nm &lt;1 ppb; 365nm &lt;1 ppb Suitability for LC-MS passes test Trace ionic impurities: Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn &lt;10 ppb each Al, Ca &lt;25 ppb each Na &lt;50 ppb Application: HPLC critical gradient applications, LC-MS, UHPLC</p>
<p>1LT H720M 2½LT H720L Dgr H:225-319-335-351-EUH019 P:210-240-305+351+338-308+313-403+233</p> 	<p><b>Tetrahydrofuran UpS</b> <b>ultra lc</b> <span style="float: right;">H720</span></p> <p><chem>CH2(CH2)2CH2O</chem> MW 72.11 BP 66.0°C d 0.89 CAS [109-99-9] Assay &gt;99.9% Water &lt;0.005% Residue &lt;0.0001% UV: 215nm &gt;10%; 235nm &gt;50%; 255nm &gt;80%; 275nm &gt;95%; 295-400nm &gt;99% Unstabilised Peroxides (at time of manufacture) &lt;0.0001% (&lt;1 ppm) Acidity &lt;0.0005 meq/g Alkalinity &lt;0.0005 meq/g Gradient Use Test: 290nm &lt;0.0005 AU Fluorescence (as quinine): 254nm &lt;1 ppb; 365nm &lt;1 ppb Suitability for LC-MS passes test Trace ionic impurities: Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn &lt;10 ppb each Al, Ca &lt;25 ppb each Na &lt;50 ppb Application: HPLC critical gradient applications, LC-MS, UHPLC</p>
<p>2½LT H772L Dgr H:225-304-315-336-361d-373 P:210-240-301+310-331-302+352-403+235</p> 	<p><b>Toluene UpS</b> <b>ultra gc</b> <span style="float: right;">H772</span></p> <p><chem>C6H5CH3</chem> MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3] Assay &gt;99.9% Water &lt;0.01% Residue &lt;0.0001% Suitability for GC-FID passes test Suitability for GC-ECD passes test Suitability for GC-MS passes test Application: GC critical low residue applications</p>
<p>2½LT H903L Dgr H:225-304-315-336-410 P:210-233-240-273-301+310-331-302+352-304+340-403+235</p> 	<p><b>2,2,4-Trimethylpentane UpS</b> <b>ultra gc</b> <span style="float: right;">H903</span></p> <p><b>(iso-Octane)</b> <chem>(CH3)3CCH2CH(CH3)2</chem> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1] Assay &gt;99.75% Water &lt;0.01% Residue &lt;0.0001% Suitability for GC-FID passes test Suitability for GC-ECD passes test Suitability for GC-MS passes test Application: GC critical low residue applications</p>

## ROMIL-UpS™ Ultra Purity Solvents Specifications

for critical analytical applications

### Water UpS

#### ultra lc

H949

1LT H949M  
2½LT H949L

H<sub>2</sub>O MW 18.02 FP 0.0°C BP 100.0°C d 1.00 CAS [7732-18-5]  
Residue <0.00005%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
TOC (at time of manufacture) <10 ppb  
Gradient Use Test: 205nm <0.002 AU; 254nm <0.0005 AU  
Baseline drift <0.02 AU @ 205nm  
Suitability for LC-MS passes test  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Filtered to 0.2 micron  
Application: HPLC critical gradient applications, LC-MS, UHPLC

### Water UpS

#### ultra pfas

H952

2½LT H952L

H<sub>2</sub>O MW 18.02 FP 0.0°C BP 100.0°C d 1.00 CAS [7732-18-5]  
Residue <0.00005%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
TOC (at time of manufacture) <10 ppb  
Gradient Use Test: 205nm <0.002 AU; 254nm <0.0005 AU  
Baseline drift <0.02 AU @ 205nm  
Suitability for LC-MS passes test  
Trace ionic impurities:  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Suitability for PFAS analysis passes test  
Filtered to 0.2 micron  
Application: Ultra low background solvent for LC-MS analysis of poly- and per-  
fluoroalkyl substances

## ROMIL Hi-Dry® Anhydrous Solvents Specifications

Acetic Acid glacial (see Acetic Acid)

### Acetic Acid Hi-Dry

D4016

100ml D4016S  
500ml D4016P  
1LT D4016M  
2½LT D4016L  
Dgr H:226-314  
P:280c-301+330+331-305+351+338-307+310

**(Acetic Acid glacial)**  
CH<sub>3</sub>COOH MW 60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7] Assay >99.8% Water <0.0050% Residue <0.0001%



### Acetone Hi-Dry

D4032

100ml D4032S  
500ml D4032P  
1LT D4032M  
2½LT D4032L  
Dgr H:225-319-336-EU066  
P:210-233-305+351+338

**(Propanone)**  
(CH<sub>3</sub>)<sub>2</sub>COMW 58.08 BP 56.1°C d 0.79 CAS [67-64-1] Assay >99.9% Water <0.0050% Residue <0.0001%



### Acetonitrile Hi-Dry

D4049

100ml D4049S  
500ml D4049P  
1LT D4049M  
2½LT D4049L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

**(Methyl Cyanide)**  
CH<sub>3</sub>CNMW41.05 BP 81.6°C d 0.78 CAS [75-05-8] Assay >99.9% Water <0.0010% Residue <0.0001%



### Acetonitrile Hi-Dry over molecular sieve

F8049

100ml F8049S  
500ml F8049P  
1LT F8049M  
2½LT F8049L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

**(Methyl Cyanide)**  
CH<sub>3</sub>CNMW41.05 BP 81.6°C d 0.78 CAS [75-05-8] Assay >99.9% Water <0.0010% Contains molecular sieve.



### Anisole Hi-Dry

D4063

100ml D4063S  
500ml D4063P  
1LT D4063M  
2½LT D4063L  
Wng H:226  
P:210-262

**(Methoxybenzene, Methyl Phenyl Ether)**  
CH<sub>3</sub>OC<sub>6</sub>H<sub>5</sub> MW108.14 BP154°C d 0.99 CAS [100-66-3] Assay >99.7% Water <0.0020% Residue <0.0005%



tert-Butanol (see 2-Methylpropan-2-ol)

n-Butanol (see Butan-1-ol)

2-Butanone (see Methyl Ethyl Ketone)

n-Butyl Alcohol (see Butan-1-ol)

tert-Butyl Alcohol (see 2-Methylpropan-2-ol)

n-Butyl Chloride (see 1-Chlorobutane)

tert-Butyl Methyl Ether (see Methyl tert-Butyl Ether)

### Butan-1-ol Hi-Dry

D4082

100ml D4082S  
500ml D4082P  
1LT D4082M  
2½LT D4082L  
Dgr H:226-302-315-318-335-336  
P:210-280F-302+352-304+340-305+351+338-313

**(n-Butanol, n-Butyl Alcohol)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>OH MW74.12 BP 117.7°C d 0.81 CAS [71-36-3] Assay >99.8% Water <0.0050% Residue <0.0005%



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### n-Butyl Acetate Hi-Dry

D4087

100ml D4087S  
500ml D4087P  
1LT D4087M  
2½LT D4087L  
Wng H:226-336-EUH066  
P:210

CH<sub>3</sub>COO(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub> MW116.16 BP126.1°C d 0.88 CAS [123-86-4] □  
Assay >99.7% Water <0.0025% Residue <0.0001%



### Carbon Disulphide Hi-Dry

D4095

100ml D4095S  
500ml D4095P  
Dgr H:225-361f-d-372-319-315  
P:210-233-280-302+352-305+351+338-403+235

CS<sub>2</sub>MW76.13 BP46.2°Cd1.26 CAS[75-15-0]  
Assay >99.9% Water <0.0050% Residue <0.0001%



### Chlorobenzene Hi-Dry

D4104

100ml D4104S  
500ml D4104P  
1LT D4104M  
2½LT D4104L  
Wng H:226-332-315-411  
P:210-273-302+352-304+340

C<sub>6</sub>H<sub>5</sub>ClMW112.56 BP131.7°C d1.11 CAS [108-90-7]  
Assay >99.9% Water <0.0020% Residue <0.0001%



### 1-Chlorobutane Hi-Dry

D4118

100ml D4118S  
500ml D4118P  
1LT D4118M  
2½LT D4118L  
Dgr H:225  
P:210

**(n-Butyl Chloride)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>Cl MW 92.57 BP 78.4°C d 0.88 CAS [109-69-3]  
Assay >99.9% Water <0.0020% Residue <0.0001%



### Chloroform Hi-Dry

stabilised with amylene

D4140

100ml D4140S  
500ml D4140P  
1LT D4140M  
2½LT D4140L  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280f-304+340-305+351+338-308+313

**(Trichloromethane)**  
CHCl<sub>3</sub> MW119.38 BP 61.2°C d 1.48 CAS [67-66-3]  
Assay >99.9%\* Water <0.0025% Residue <0.0001%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm



### Chloroform Hi-Dry

stabilised with amylene over molecular sieve

F8140

1 00ml F81 40S  
500ml F8140P  
1LT F8140M  
2½LT F8140L  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280f-304+340-305+351+338-308+313

**(Trichloromethane)**  
CHCl<sub>3</sub> MW119.38 BP 61.2°C d 1.48 CAS [67-66-3]  
Assay >99.9%\* Water <0.0025%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm  
Contains molecular sieve.



### Cyclohexane Hi-Dry

D4156

100ml D4156S  
500ml D4156P  
1LT D4156M  
2½LT D4156L  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-302+352-331-403+235

C<sub>6</sub>H<sub>12</sub> MW84.16FP 6.5°C BP 80.7°C d 0.78 CAS [110-82-7]  
Assay >99.9% Water <0.0005% Residue <0.0001%

## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### Cyclohexane Hi-Dry

over molecular sieve

F8156

100ml F8156S  
500ml F8156P  
1LT F8156M  
2½LT F8156L  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-  
302+352-331-403+235



C6H12 MW 84.16 FP 6.5°C BP 80.7°C d 0.78 CAS [110-82-7]

Assay >99.9% Water  
<0.0005% Contains molecular  
sieve.

### Cyclohexanone Hi-Dry

D4173

100ml D4173S  
500ml D4173P  
1LT D4173M  
2½LT D4173L  
Wng H:226-332  
P:210



C6H10O MW 98.15 FP -47°C BP 155°C d 0.94 CAS [108-94-  
1] Assay >99.8% Water <0.0050% Residue <0.0005%

### 1,2-Dichlorobenzene Hi-Dry

D4178

100ml D4178S  
500ml D4178P  
1LT D4178M  
2½LT D4178L  
Wng H:302-315-319-335-410  
P:273-302+352-  
305+351+338



C6H4Cl2MW 147.00 FP-17°C BP180.5°Cd1.31 CAS [95-50-1]  
Assay >99.8% Water <0.0020% Residue <0.0005%

### Dichloromethane Hi-Dry

stabilised with amylene

D4202

100ml D4202S  
500ml D4202P  
1LT D4202M  
2½LT D4202L  
Wng H:351  
P:281-308+313



(Methylene Dichloride)  
CH2Cl2MW84.93BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.0020% Residue <0.0001%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm

### Dichloromethane Hi-Dry

stabilised with amylene over molecular sieve

F8202

100ml F8202S  
500ml F8202P  
1LT F8202M  
2½LT F8202L  
Wng H:351  
P:281-308+313



(Methylene Dichloride)  
CH2Cl2MW84.93BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.0020%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm  
Contains molecular sieve.

### Dichloromethane Hi-Dry

stabilised with amylene extra dry

D4203

100ml D4203S  
500ml D4203P  
1LT D4203M  
2½LT D4203L  
Wng H:351  
P:281-308+313



(Methylene Dichloride)  
CH2Cl2MW84.93BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.0010% Residue <0.0001%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm

### Diethyl Ether Hi-Dry

stabilised with ethanol

D4219

100ml D4219S  
500ml D4219P  
1LT D4219M  
2½LT D4219L  
Dgr H:224-302-336-EUH019-EUH066  
P:210-240-403+235



(C2H5)2O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-  
7] Assay >99.9%\* Water <0.0025% Residue  
<0.0001% \*ex stabiliser  
Stabiliser: Ethanol ca. 1% v/v



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### Diethyl Ether Hi-Dry

stabilised with ethanol over molecular sieve

F8219

100ml F8219S (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7] Assay >99.9%\* Water <0.0025%  
 500ml F8219P  
 1LT F8219M  
 2½LT F8219L  
 Dgr H:224-302-336-EUH019-EUH066  
 P:210-240-403+235



### Diethyl Ether Hi-Dry

stabilised with BHT

D4220

100ml D4220S (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7] Assay >99.9%\* Water <0.0025% Residue <0.0001%\*  
 500ml D4220P  
 1LT D4220M  
 2½LT D4220L  
 Dgr H:224-302-336-EUH019-EUH066  
 P:210-240-403+235



### Diethyl Ether Hi-Dry

stabilised with BHT over molecular sieve

F8220

100ml F8220S (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7] Assay >99.9%\* Water <0.0025%  
 500ml F8220P  
 1LT F8220M  
 2½LT F8220L  
 Dgr H:224-302-336-EUH019-EUH066  
 P:210-240-403+235



### 1,2-Dimethoxyethane Hi-Dry

D4261

100ml D4261S (Ethylene Glycol Dimethyl Ether)  
 500ml D4261P CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> MW90.12 FP -58°C BP 85°C d 0.87 CAS [110-71-4] Assay >99.8% Water <0.0030% Residue <0.0005%  
 1LT D4261M  
 2½LT D4261L  
 Dgr H:225-360FD-332-EUH019 P:201-210-308+313-403+235



### 1,2-Dimethoxyethane Hi-Dry

over molecular sieve

F8261

100ml F8261S (Ethylene Glycol Dimethyl Ether)  
 500ml F8261P CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> MW90.12 FP -58°C BP 85°C d 0.87 CAS [110-71-4] Assay >99.8% Water <0.0030%  
 1LT F8261M  
 2½LT F8261L  
 Dgr H:225-360FD-332-EUH019 P:201-210-308+313-403+235



### Dimethylacetamide Hi-Dry

D4248

100ml D4248S CH<sub>3</sub>CON(CH<sub>3</sub>)<sub>2</sub> MW 87.12 FP -20°C BP 166.1°C d 0.94 CAS [127-19-5] Assay >99.7% Water <0.0050% Residue <0.0005%  
 500ml D4248P  
 1LT D4248M  
 2½LT D4248L  
 Dgr H:227-360D-312+332-319  
 P:201-302+352-305+351+338-308+313



### Dimethylformamide Hi-Dry

over molecular sieve

F8252

100ml F8252S HCON(CH<sub>3</sub>)<sub>2</sub> MW 73.09 BP 153.0°C d 0.95 CAS [68-12-2] Assay >99.9% Water <0.0050%  
 500ml F8252P  
 1LT F8252M  
 2½LT F8252L  
 Dgr H:360D-226-312+332-319  
 P:201-210-302+352-305+351+338-308+313



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### 1,4-Dioxan Hi-Dry

D4297

100ml D4297S  
500ml D4297P  
1LT D4297M  
2½LT D4297L  
Dgr H:225-350-319-335-EUH019-  
EUH066  
P:210-281-305+351+338-308+313

C4H8O2 MW 88.11 FP 11.8°C BP 101.3°C d 1.03 CAS [123-91-1] Assay >99.9% Water <0.0025% Residue <0.0001% Unstabilised



### 1,4-Dioxan Hi-Dry

over molecular sieve

F8297

100ml F8297S  
500ml F8297P  
1LT F8297M  
2½LT F8297L  
Dgr H:225-350-319-335-EUH019-  
EUH066  
P:210-281-305+351+338-308+313

C4H8O2 MW 88.11 FP 11.8°C BP 101.3°C d 1.03 CAS [123-91-1] Assay >99.9% Water <0.0025% Unstabilised  
Contains molecular sieve.



### Di-iso-propyl Ether Hi-Dry

stabilised with BHT

D4236

100ml D4236S  
500ml D4236P  
1LT D4236M  
2½LT D4236L  
Dgr H:225-336-EUH019-EUH066  
P:210-240-403+235

[(CH3)2CH]2O MW 102.18 BP 68.5°C d 0.73 CAS [108-20-3] Assay >99.5%\* Water <0.0025% Residue <0.0001%\*  
\*ex stabiliser  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 5 ppm



### Di-iso-propylethylamine Hi-Dry

D4240

100ml D4240S  
500ml D4240P  
Dgr H:225-301-314-412  
P:210-233-240-273-280-  
301+330+331-305+351+338-  
309+310-403+235

(Ethyl-di-iso-propylamine)  
C8H19NMW 129.25 BP 127°C d 0.76 CAS [7087-68-5] Assay >99.5% Water <0.0100% Residue <0.0001%



Ethyl Alcohol (see Ethanol)

Ethyl-di-iso-propylamine (see Di-iso-propylethylamine)

Ethylene Glycol (see 1,2-Ethanediol)

Ethylene Glycol Dimethyl Ether (see 1,2-Dimethoxyethane)

### 1,2-Ethanediol Hi-Dry

D4348

100ml D4348S  
500ml D4348P  
1LT D4348M  
2½LT D4348L  
Wng H:302

(Ethylene Glycol)  
CH2(OH)CH2OH MW 62.07 FP -13°C BP 197.3°C CAS [107-21-1] Assay >99.8% Water <0.0100% Residue <0.0005%



### Ethanol absolute Hi-Dry

D4313

100ml D4313S  
500ml D4313P  
1LT D4313M  
2½LT D4313L  
Dgr H:225  
P:210-233-240-403+235

(Ethyl Alcohol)  
C2H5OHMW 46.07 BP 78.3°C d 0.79 CAS [64-17-5] Assay >99.8% Water <0.0050% Residue <0.0005%



### Ethyl Acetate Hi-Dry

D4346

100ml D4346S  
500ml D4346P  
1LT D4346M  
2½LT D4346L  
Dgr H:225-319-336-EUH066  
P:210-233-240-305+351+338-  
403+235

CH3COOC2H5 MW 88.11BP77.1°C d 0.90 CAS [141-78-6] Assay >99.9% Water <0.0025% Residue <0.0001%



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### Ethyl Acetate Hi-Dry

over molecular sieve

F8346

100ml F8346S CH<sub>3</sub>COOC<sub>2</sub>H<sub>5</sub> MW 88.11 BP 77.1°C d 0.90 CAS [141-78-6] Assay >99.9% Water <0.0025%  
 500ml F8346P  
 1LT F8346M  
 2½LT F8346L  
 Dgr H:225-319-336-EU066  
 P:210-233-240-305+351+338-403+235



### Heptane fraction Hi-Dry

D4368

100ml D4368S C<sub>7</sub>H<sub>16</sub> BP 85-99°C d 0.69  
 500ml D4368P Water <0.0005% Residue <0.0001%  
 1LT D4368M Comprises ca. 20-50% n-isomer, the remainder being predominantly other  
 2½LT D4368L isomers of heptane.  
 Dgr H:225-304-315-336-410  
 P:210-273-301+310-331-302+352-304+340-403+235



### n-Heptane 95% Hi-Dry

D4367

100ml D4367S CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW100.21BP94-98°C d 0.68 CAS [142-82-5]  
 500ml D4367P Water <0.0005% Residue <0.0001%  
 1LT D4367M Assay (n-isomer) >95%  
 2½LT D4367L Assay (all isomers) >99.5%  
 Dgr H:225-304-315-336-410  
 P:210-273-301+310-331-302+352-304+340-403+235



### n-Heptane 95% Hi-Dry

over molecular sieve

F8367

100ml F8367S CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW 100.21BP 94-98°C d 0.68 CAS [142-82-5]  
 500ml F8367P Water <0.0005%  
 1LT F8367M Assay (n-isomer) >95%  
 2½LT F8367L Assay (all isomers) >99.5%  
 Dgr H:225-304-315-336-410  
 P:210-273-301+310-331-302+352-304+340-403+235



### n-Heptane 99% Hi-Dry

D4366

100ml D4366S CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW 100.21BP 98.4°C d 0.68 CAS [142-82-5]  
 500ml D4366P Assay >99% Water <0.0005% Residue <0.0001%  
 1LT D4366M  
 2½LT D4366L  
 Dgr H:225-304-315-336-410  
 P:210-273-301+310-331-302+352-304+340-403+235



### Hexane fraction Hi-Dry

D4390

100ml D4390S C<sub>6</sub>H<sub>14</sub> BP 65-70°C d 0.66 CAS [73513-42-5]  
 500ml D4390P Water <0.0005% Residue <0.0001%  
 1LT D4390M Comprises ca. 50% n-isomer, the remainder being predominantly other  
 2½LT D4390L isomers of hexane.  
 Dgr H:225-304-361F-373-315-336-411  
 P:210-240-273-301+310-331-302+352-403+235



### iso-Hexane 95% Hi-Dry

D4388

100ml D4388S C<sub>6</sub>H<sub>14</sub> MW 86.18 BP55-63°C d 0.65CAS [107-83-5]  
 500ml D4388P Water <0.0005% Residue <0.0001%  
 1LT D4388M n-Hexane < 5%  
 2½LT D4388L  
 Dgr H:225-304-315-336-411  
 P:233-273-301+310-331-302+352-403+235



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### n-Hexane 95% Hi-Dry

D4389

100ml D4389S  
500ml D4389P  
1LT D4389M  
2½LT D4389L  
Dgr H:225-304-361f-373-315-336-411  
P:210-240-273-301+310-331-302+352-403+235

CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> MW 86.18 BP 67-70°C d 0.66 CAS [110-54-3]  
Water <0.0005% Residue <0.0001%  
Assay (n-isomer) >95%  
Assay (all isomers) >99.5%



### n-Hexane 95% Hi-Dry

over molecular sieve

F8389

100ml F8389S  
500ml F8389P  
1LT F8389M  
2½LT F8389L  
Dgr H:225-304-361f-373-315-336-411  
P:210-240-273-301+310-331-302+352-403+235

CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> MW 86.18 BP 67-70°C d 0.66 CAS [110-54-3]  
Water <0.0005%  
Assay (n-isomer) >95%  
Assay (all isomers) >99.5%  
Contains molecular sieve.



Methoxybenzene (see Anisole)

Methyl Alcohol (see Methanol)

Methyl Cyanide (see Acetonitrile)

4-Methyl-1,3-dioxolan-2-one (see Propylene Carbonate)

Methylene Dichloride (see Dichloromethane)

4-Methylpentan-2-one (see Methyl iso-Butyl Ketone)

Methyl Phenyl Ether (see Anisole)

### Methanol Hi-Dry

D4412

100ml D4412S  
500ml D4412P  
1LT D4412M  
2½LT D4412L  
Dgr H:225-301+311+331-370  
P:210-280f-302+352-309+310-403+235

(Methyl Alcohol)  
CH<sub>3</sub>OH MW32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
Assay >99.9% Water <0.0035% Residue <0.0001%



### Methanol Hi-Dry

over molecular sieve

F8412

100ml F8412S  
500ml F8412P  
1LT F8412M  
2½LT F8412L  
Dgr H:225-301+311+331-370  
P:210-280f-302+352-309+310-403+235

(Methyl Alcohol)  
CH<sub>3</sub>OH MW32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
Assay >99.9% Water <0.0035%  
Contains molecular sieve.



### bis(2-Methoxyethyl) Ether Hi-Dry

D4068

100ml D4068S  
500ml D4068P  
1LT D4068M  
Dgr H:226-360FD-EUH019  
P:201-210-308+313

(Diglyme, Diethylene Glycol Dimethyl Ether)  
(CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>OMW134.17 BP162°C d 0.94 CAS [111-96-6]  
Assay >99.8% Water <0.0030% Residue <0.0005%  
Unstabilised



### bis(2-Methoxyethyl) Ether Hi-Dry

over molecular sieve

F8068

100ml F8068S  
500ml F8068P  
1LT F8068M  
Dgr H:226-360FD-EUH019  
P:201-210-308+313

(Diglyme, Diethylene Glycol Dimethyl Ether)  
(CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>OMW134.17BP162°C d 0.94 CAS [111-96-6]  
Assay >99.8% Water <0.0020%  
Unstabilised  
Contains molecular sieve.



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### Methyl tert-Butyl Ether Hi-Dry

D4447

100ml D4447S  
500ml D4447P  
1LT D4447M  
2½LT D4447L  
Dgr H:225-315  
P:210-233-302+352-403+235

(tert-Butyl Methyl Ether)

CH<sub>3</sub>OC(CH<sub>3</sub>)<sub>3</sub> MW88.15 BP 55.4°C d 0.74 CAS [1634-04-4] Assay >99.7% Water <0.0050% Residue <0.0001%



### Methyl iso-Butyl Ketone Hi-Dry

D4445

100ml D4445S  
500ml D4445P  
1LT D4445M  
2½LT D4445L  
Dgr H:225-332-319-336-351-EUH066  
P:210-305+351+338-304+340

(4-Methylpentan-2-one)

(CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>COCH<sub>3</sub>MW 100.16 BP 117.4°C d 0.80 CAS [108-10-1] Assay >99.7% Water <0.0050% Residue <0.0005%



### Methylcyclohexane Hi-Dry

D4465

100ml D4465S  
500ml D4465P  
Dgr H:225-304-315-336-411  
P:273-301+310-331-302+352-403+235

C<sub>7</sub>H<sub>14</sub> MW 98.19 BP 101°C d 0.77 CAS [108-87-2]

Assay >99.9% Water <0.0005% Residue <0.0001%



### Methylcyclopentane 95% Hi-Dry

D4473

100ml D4473S  
500ml D4473P  
1LT D4473M  
Dgr H:225-304  
P:210-260v-262-301+310-331-403+235

C<sub>6</sub>H<sub>12</sub> MW 84.16 BP 69-73°C d 0.75 CAS [96-37-7]

Water <0.0005% Residue <0.0001%

Comprises ca. 95% methylcyclopentane, the remainder being predominantly other C<sub>6</sub>H<sub>12</sub> isomers.



### Methyl Ethyl Ketone Hi-Dry

D4494

100ml D4494S  
500ml D4494P  
1LT D4494M  
2½LT D4494L  
Dgr H:225-319-336-EUH066  
P:210-305+351+338-403+233

(2-Butanone)

CH<sub>3</sub>CH<sub>2</sub>COCH<sub>3</sub> MW 72.11 BP 79.6°C d 0.80 CAS [78-93-3] Assay >99.8% Water <0.0050% Residue <0.0001%



### N-Methyl-2-pyrrolidone Hi-Dry

D4565

100ml D4565S  
500ml D4565P  
1LT D4565M  
2½LT D4565L  
Dgr H:360D-315-319-335  
P:201-302+352-305+351+338-308+313

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONCH<sub>3</sub> MW 99.13 BP 202.0°C d 1.03 CAS [872-50-4] Assay >99.5% Water <0.0075%



### N-Methyl-2-pyrrolidone Hi-Dry

F8564

100ml F8564S  
500ml F8564P  
1LT F8564M  
2½LT F8564L  
Dgr H:360D-315-319-335  
P:201-302+352-305+351+338-308+313

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONCH<sub>3</sub> MW 99.13 BP 202.0°C d 1.03 CAS [872-50-4] Assay >99.5% Water <0.0050%

Contains molecular sieve.



### 2-Methyltetrahydrofuran Hi-Dry

D4536

100ml D4536S  
500ml D4536P  
1LT D4536M  
2½LT D4536L  
Dgr H:225-319-335-EUH019  
P:210-233-240-305+351+338-403+235

CH<sub>3</sub>C<sub>4</sub>H<sub>7</sub>O MW 86.13 BP 80°C d 0.86 CAS [96-47-9]

Assay >99.8% Water <0.0025% Residue <0.0001%

Unstab liised






## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### 2-Methyltetrahydrofuran Hi-Dry stabilised with BHT


D4537

100ml D4537S CH<sub>3</sub>C<sub>4</sub>H<sub>7</sub>O MW 86.13 BP 80°C d 0.86 CAS [96-47-9]   
 500ml D4537P Assay >99.8%\* Water <0.0025% Residue <0.0001%\*  
 1LT D4537M \*ex stabiliser  
 2½LT D4537L Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm  
 Dgr H:225-319-335-EUH019  
 P:210-233-240-305+351+338-403+235



### n-Nonane 95% Hi-Dry

D4568

100ml D4568S CH<sub>3</sub>(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub> MW128.26BP146-150°C d 0.72 CAS [111-84-2]   
 500ml D4568P Water <0.0005% Residue <0.0001%  
 1LT D4568M  
 2½LT D4568L  
 Dgr H:226-304  
 P:210-301+310-331



### iso-Octane (see 2,2,4-Trimethylpentane)

Perchloroethylene (see Tetrachloroethylene)

Petroleum Distillate (see Petroleum Ether)

Petroleum Spirit (see Petroleum Ether)

n-Propanol (see Propan-1-ol)

iso-Propanol (see Propan-2-ol)


Propanone (see Acetone)

n-Propyl Alcohol (see Propan-1-ol)

iso-Propyl Alcohol (see Propan-2-ol)

### n-Pentane 95% Hi-Dry


D4571

100ml D4571S CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub> MW72.15 BP35.5-37°C d 0.63 CAS [109-66-0]   
 500ml D4571P Water <0.0005% Residue <0.0001%  
 1LT D4571M Assay (n-isomer) >95%  
 2½LT D4571L Assay (all isomers) >99.5%  
 Dgr H:225-304-336-411-EUH066  
 P:273-301+310-331-403+235



### Petroleum Ether 30-40°C Hi-Dry


D4600

100ml D4600S (Petroleum Distillate, Petroleum Spirit)   
 500ml D4600P BP 30-40°C d0.62 CAS[109-66-0]  
 1LT D4600M Water <0.0005% Residue <0.0001%  
 2½LT D4600L  
 Dgr H:225-304-336-411-EUH066  
 P:210-243-301+310-303+361+353-405-501



### Petroleum Ether 40-60°C Hi-Dry


D4601

100ml D4601S (Petroleum Distillate, Petroleum Spirit)   
 500ml D4601P BP 40-60°C d0.64 CAS[8032-32-4]  
 1LT D4601M Water <0.0005% Residue <0.0001%  
 2½LT D4601L  
 Dgr H:225-304-336-411-EUH066  
 P:210-233-243-273-280-301+310-303+361+353-304-331-403+235










### Petroleum Ether 60-80°C Hi-Dry

D4602

100ml D4602S (Petroleum Distillate, Petroleum Spirit)   
 500ml D4602P BP 60-80°C d0.67  
 1LT D4602M Water <0.0005% Residue <0.0001%  
 2½LT D4602L  
 Dgr H:225-304-315-336-411  
 P:210-243-273-280-301+310-331-403+235



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

100ml D4605S 500ml D4603P 1LT D4605M 2½LT D4605L Dgr H:225-304-315-336-411 P:210-273-280-260v+301+310-331-403+235	<b>Petroleum Ether 80-100°C Hi-Dry</b> <b>(Petroleum Distillate, Petroleum Spirit)</b> BP80-100°C d 0.69 CAS [64742-49-0] Water <0.0005% Residue <0.0001%	<b>D4603</b>	
	<b>Propan-1-ol Hi-Dry</b> <b>(n-Propanol, n-Propyl Alcohol)</b> CH3CH2CH2OH MW 60.10 BP 97.2°C d 0.80 CAS [71-23-8] Assay >99.9% Water <0.0050% Residue <0.0005%	<b>D4623</b>	
100ml D4625S 500ml D4625P 1LT D4625M 2½LT D4625L Dgr H:225-318-336 P:210-233-280f-305+351+338-313		<b>Propan-2-ol Hi-Dry</b> <b>(iso-Propanol, iso-Propyl Alcohol)</b> (CH3)2CHOH MW 60.10 BP 82.2°C d 0.78 CAS [67-63-0] Assay >99.9% Water <0.0050% Residue <0.0005%	<b>D4625</b>
100ml D4625S 500ml D4625P 1LT D4625M 2½LT D4625L Dgr H:225-319-336 P:210-233-305+351+338		<b>Propan-2-ol Hi-Dry</b> <b>over molecular sieve</b> <b>(iso-Propanol, iso-Propyl Alcohol)</b> (CH3)2CHOH MW 60.10 BP 82.2°C d 0.78 CAS [67-63-0] Assay >99.9% Water <0.0050% Contains molecular sieve.	<b>F8625</b>
100ml F8625S 500ml F8625P 1LT F8625M 2½LT F8625L Dgr H:225-319-336 P:210-233-305+351+338		<b>Propylene Carbonate Hi-Dry</b> <b>(4-Methyl-1,3-dioxolan-2-one)</b> CH3CHOCOOCH2 MW 102.09 BP 241.7°C d 1.20 CAS [108-32-7] Assay >99.8% Water <0.0050% Residue <0.0001%	<b>D4645</b>
100ml D4645S 500ml D4645P 1LT D4645M Wng H:319 P:305+351+338		<b>Pyridine Hi-Dry</b> <b>over molecular sieve</b> C5H5N MW 79.10 BP 115.3°C d 0.98 CAS [110-86-1] Assay >99.8% Water <0.0050% Contains molecular sieve.	<b>F8652</b>
100ml F8652S 500ml F8652P 1LT F8652M 2½LT F8652L Dgr H:225-302+312+332-315-319 P:210-302+352-304+340-305+351+338-403+235		<b>Tetramethylene Sulphone (see Sulpholane)</b> <b>Trichloromethane (see Chloroform)</b>	
100ml D4702S 500ml D4702P 1LT D4702M 2½LT D4702L Wng H:315-317-319-336-351-411 P:273-281-302+352-305+351+338-308+313		<b>Tetrachloroethylene Hi-Dry</b> <b>(Perchloroethylene)</b> CCl2CCl2 MW 165.83 BP 121.1°C d 1.62 CAS [127-18-4] Assay >99.9% Water <0.0020% Residue <0.0001% Unstabilised	<b>D4702</b>

## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### Tetrahydrofuran Hi-Dry

D4718

100ml D4718S  
500ml D4718P  
1LT D4718M  
2½LT D4718L  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66°C d 0.89 CAS [109-99-9]  
Assay >99.9% Water <0.0025% Residue <0.0001%  
Unstabilised



### Tetrahydrofuran Hi-Dry

over molecular sieve

F8717

100ml F8717S  
500ml F8717P  
1LT F8717M  
2½LT F8717L  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66°C d 0.89 CAS [109-99-9]  
Assay >99.9% Water <0.0025%  
Unstabilised  
Contains molecular sieve.



### Tetrahydrofuran Hi-Dry

stabilised with BHT

D4719

100ml D4719S  
500ml D4719P  
1LT D4719M  
2½LT D4719L  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66°C d 0.89 CAS [109-99-9]  
Assay >99.9%\* Water <0.0025% Residue <0.0001%\*  
\*ex stabiliser  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm



### Toluene Hi-Dry

D4771

100ml D4771S  
500ml D4771P  
1LT D4771M  
2½LT D4771L  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235

C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub> MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]  
Assay >99.9% Water <0.0010% Residue <0.0001%



### Toluene Hi-Dry

over molecular sieve

F8771

100ml F8771S  
500ml F8771P  
1LT F8771M  
2½LT F8771L  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235

C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub> MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]  
Assay >99.9% Water <0.0010%  
Contains molecular sieve.



### Triethylamine Hi-Dry

D4763

100ml D4763S  
500ml D4763P  
1LT D4763M  
Dgr H:225-302+312+332-314-335  
P:210-280-301+330+331-302+352-304+340-305+351+338-309+310-403+235

(CH<sub>3</sub>CH<sub>2</sub>)<sub>3</sub>N MW 101.19 BP 88.9°C d 0.73 CAS [121-44-8]  
Assay >99.8% Water <0.0050% Residue <0.0005%



### Triethylamine Hi-Dry

over molecular sieve

F8763

100ml F8763S  
500ml F8763P  
1LT F8763M  
Dgr H:225-302+312+332-314-335  
P:210-280-301+330+331-302+352-304+340-305+351+338-309+310-403+235

(CH<sub>3</sub>CH<sub>2</sub>)<sub>3</sub>N MW 101.19 BP 88.9°C d 0.73 CAS [121-44-8]  
Assay >99.8% Water <0.0050%  
Contains molecular sieve.



## ROMIL Hi-Dry® Anhydrous Solvents Specifications

### 2,2,4-Trimethylpentane Hi-Dry

D4901

100ml D4901S (iso-Octane) □  
 500ml D4901P (CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1] Assay >99.75% Water <0.0005% Residue <0.0001%  
 1LT D4901M  
 2½LT D4901L  
 Dgr H:225-304-315-336-410  
 P:210-233-240-273-301+310-331-302+352-304+340-403+235



### 2,2,4-Trimethylpentane Hi-Dry

over molecular sieve

F8901

100ml F8901S (iso-Octane) □  
 500ml F8901P (CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1] Assay >99.75% Water <0.0005%  
 1LT F8901M  
 2½LT F8901L  
 Dgr H:225-304-315-336-410  
 P:210-233-240-273-301+310-331-302+352-304+340-403+235  
 Contains molecular sieve.



### Xylene mixed isomers Hi-Dry

D4982

100ml D4982S C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW 106.17 BP 138-142°C d 0.86 CAS [1330-20-7] □  
 500ml D4982P Water <0.0010% Residue <0.0001%  
 1LT D4982M Comprises 3 isomers and ethylbenzene  
 2½LT D4982L Assay (total C<sub>8</sub>H<sub>10</sub> isomers) >98.5%  
 Wng H:226-312+332- Ethylbenzene typically <3%  
 315 P:210-302+352- Toluene typically <0.5%  
 304+340 Methylethylbenzene typically <0.5%



### Xylene mixed isomers Hi-Dry

over molecular sieve

F8982

100ml F8982S C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW 106.17 BP 138-142°C d 0.86 CAS [1330-20-7] Water <0.0010% □  
 500ml F8982P Comprises 3 isomers and ethylbenzene  
 1LT F8982M Assay (total C<sub>8</sub>H<sub>10</sub> isomers) >98.5%  
 2½LT F8982L Ethylbenzene typically <3%  
 Wng H:226-312+332- Toluene typically <0.5%  
 315 P:210-302+352- Methylethylbenzene typically <0.5%  
 304+340 Contains molecular sieve.



## ROMIL Hi-Dry® KF Karl Fischer Reagents and Calibrants Specifications

pyridine-free reagents for water determination

### Benzoic Acid Hi-Dry KF

K8114

500g K8114P  
Wng H:302-319  
P:305+351+338

C6H5COOH MW122.12CAS [65-85-0]  
Water <0.2%  
Application: Buffer reagent for KF titration



### Chloroform Hi-Dry KF

stabilised with amylene

K8140

1LT K8140M  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280f-304+340-305+351+338-308+313

(Trichloromethane)  
CHCl<sub>3</sub> MW 119.38 BP 61.2°C d 1.48 CAS [67-66-3]  
Water <0.0100%  
Stabiliser: Amylene ca. 25 ppm  
Application: Solvent for KF sample dissolution



### Formamide Hi-Dry KF

K8351

1LT K8351M  
Dgr H:360D  
P:201-308+313

HCONH<sub>2</sub> MW 45.04FP2.5°CBP210.5°C d 1.13 CAS [75-12-7]  
Water <0.0200%  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Store in dark.  
Application: Solvent for KF sample dissolution



### Karl Fischer electrolyte A Hi-Dry KF

K3035

500ml K3035P  
1LT K3035M  
Dgr H:225-301+311+331-314-360D-370-373  
P:201-210-280-302+352-309+310-403+235

Use in conjunction with Karl Fischer electrolyte C (ROMIL K3146) for cells with diaphragm.  
Protect from atmospheric moisture.  
Application: Pyridine-free reagent for coulometric KF titration without diaphragm and anode reagent for cells with diaphragm



### Karl Fischer electrolyte AF Hi-Dry KF

for oils and fats

K3072

100ml K3072S  
500ml K3072P  
Dgr H:225-301+311+331-314-351-360D-370-372  
P:201-210-260-280-305+351+338-308+313

Use in conjunction with Karl Fischer electrolyte C (ROMIL K3146).  
Protect from atmospheric moisture.  
Application: Pyridine-free anode reagent (chloroform/xylene-based) for coulometric KF titration of refined and crude oils and fats



### Karl Fischer electrolyte AH Hi-Dry KF

for long-chain hydrocarbons

K3041

500ml K3041P  
Dgr H:225-301+311+331-314-360D-335-370-411  
P:201-210-273-280-302+352-309+310-403+235

Use in conjunction with Karl Fischer electrolyte C (ROMIL K3146) for cells with diaphragm.  
Protect from atmospheric moisture.  
Application: Pyridine-free reagent for coulometric KF titration of long-chain hydrocarbons without diaphragm and anode reagent with diaphragm



### Karl Fischer electrolyte AK Hi-Dry KF

for aldehydes & ketones

K3068

500ml K3068P  
Dgr H:226-302-331-315-318-336-351-360FD-372  
P:201-210-280-301+330+331-302+352-304+340-305+351+338-308+313

Use in conjunction with Karl Fischer electrolyte CK (ROMIL K3179).  
Protect from atmospheric moisture.  
Application: Pyridine-free anode reagent for coulometric KF titration of aldehydes and ketones



### Karl Fischer electrolyte AO Hi-Dry KF

K3054

500ml K3054P  
Dgr H:225-301+311+331-314-360D-370-373  
P:201-210-280-302+352-309+310-403+235









Use in conjunction with Karl Fischer electrolyte C (ROMIL K3146) for cells with diaphragm.  
Protect from atmospheric moisture.  
Application: Pyridine-free reagent for coulometric KF titration with oven without diaphragm and anode reagent with diaphragm





## ROMIL Hi-Dry® KF Karl Fischer Reagents and Calibrants Specifications

pyridine-free reagents for water determination

500ml K3087P Dgr H:225-301+311+331-314-351-360D-370-372 P:201-210-280-304+340-305+351+338-308+313-403+235	<b>Karl Fischer electrolyte AP Hi-Dry KF</b> <span style="float: right;">K3087</span> Use in conjunction with Karl Fischer electrolyte C (ROMIL K3146). Protect from atmospheric moisture. Application: Pyridine-free anode reagent for coulometric KF titration with diaphragm
	<b>Karl Fischer electrolyte AR Hi-Dry KF</b> <span style="float: right;">K3029</span>
500ml K5029P Dgr H:225-301+311+331-318-360D-370-373 P:201-210-260-280-301+310-305+351+338	Protect from atmospheric moisture. Application: Pyridine-free reagent for coulometric KF titration without diaphragm
	<b>Karl Fischer electrolyte C Hi-Dry KF</b> <span style="float: right;">K3146</span>
25ml K3146V 50ml K3146T 10x5ml K3146Q Dgr H:225-301+311+331-315-318-370-373 P:210-280-302+352-305+351+338-309+310-403+235	Use in conjunction with suitable Karl Fischer electrolyte A* for cells with diaphragm. Protect from atmospheric moisture. Application: Pyridine-free cathode reagent for coulometric KF titration with diaphragm
	<b>Karl Fischer electrolyte CK Hi-Dry KF</b> <span style="float: right;">K3179</span> <b>for aldehydes &amp; ketones</b>
50ml K3179T Dgr H:312-314-360Df P:201-280-302+352-305+351+338-308+313	Use in conjunction with Karl Fischer electrolyte AK (ROMIL K3068). Protect from atmospheric moisture. Application: Pyridine-free cathode reagent for coulometric KF titration of aldehydes and ketones
	<b>Karl Fischer reagent VC2 Hi-Dry KF</b> <span style="float: right;">K5102</span> <b>pyridine-free composite</b>
1LT K5102M 2½LT K5102L Dgr H:314-332-360D-401 P:201-273-280c-304+340-305+351+338-308+310	Nominal titre 1.0 ml = 2.0 mg ±5% H <sub>2</sub> O @ 20°C Calibrate before each use. Application: Pyridine-free ready to use reagent for volumetric KF titration
	<b>Karl Fischer reagent VC5 Hi-Dry KF</b> <span style="float: right;">K5105</span> <b>pyridine-free composite</b>
500ml K5105P 1LT K5105M 2½LT K5105L Dgr H:314-332-360D-401 P:201-273-280c-304+340-305+351+338-308+310	Nominal titre 1.0 ml = 5.0 mg ±5% H <sub>2</sub> O @ 20°C Calibrate before each use. Application: Pyridine-free ready to use reagent for volumetric KF titration
	<b>Karl Fischer reagent VCK5 Hi-Dry KF</b> <span style="float: right;">K5235</span> <b>pyridine-free composite for aldehydes &amp; ketones</b>
1LT K5235M Dgr H:314-332-360D-401 P:201-273-280c-304+340-305+351+338-308+310	Nominal titre 1.0 ml = 5.0 mg ±5% H <sub>2</sub> O @ 20°C Calibrate before each use. Application: Pyridine-free ready to use reagent for volumetric KF titration of aldehydes and ketones
	<b>Karl Fischer solvent VK Hi-Dry KF</b> <span style="float: right;">K5246</span> <b>for aldehydes &amp; ketones</b>
1LT K5246M Dgr H:300+310+330-351-361d-372-315-319 P:201-260v-280f-302+350-304+340-305+351+338-308+313	Use in conjunction with Karl Fischer reagent VCK5 (ROMIL K5235). Protect from atmospheric moisture. Application: Solvent for volumetric KF titration of aldehydes and ketones
	

## ROMIL Hi-Dry® KF Karl Fischer Reagents and Calibrants Specifications

pyridine-free reagents for water determination

### Karl Fischer solvent VKM Hi-Dry KF

for aldehydes & ketones

K5257

1LT K5257M  
Dgr H:226-302-331-315-318-336-  
351-360D-372  
P:201-210-280-304+340-  
305+351+338-308+313-403+233



Use in conjunction with Karl Fischer reagent VCK5 (ROMIL K5235).  
Protect from atmospheric moisture.  
Application: Solvent for volumetric KF titration of aldehydes and ketones

### Karl Fischer solvent VKS Hi-Dry KF

for aldehydes & ketones

K5268

500ml K5268P  
1LT K5268M  
Dgr H:226-315-318-360D-336  
P:201-210-280-304+340-  
305+351+338-308+313



Use in conjunction with Karl Fischer reagent VCK5 (ROMIL K5235).  
Protect from atmospheric moisture.  
Application: Solvent (halogenated hydrocarbon-free) for volumetric KF titration of aldehydes and ketones

### Karl Fischer solvent VOC Hi-Dry KF

for oils and fats

K5373

1LT K5373M  
Dgr H:226-302-312-331-315-319-  
351-361d-370-372  
P:201-210-261v-280f-304+340-  
305+351+338-308+313



Use in conjunction with Karl Fischer reagent VC? (ROMIL K510?).  
Protect from atmospheric moisture.  
Application: Solvent (chloroform-based) for volumetric KF titration of oils and fats

### Karl Fischer solvent VOH Hi-Dry KF

for oils and fats

K5352

1LT K5352M  
Dgr H:225-301+311+331-319-370  
P:210-280f-302+352-309+310-  
403+235



Use in conjunction with Karl Fischer reagent VC? (ROMIL K510?).  
Protect from atmospheric moisture.  
Application: Solvent (hexan-1-ol-based) for volumetric KF titration of oils and fats

### Karl Fischer solvent VOX Hi-Dry KF

for crude oils

K5331

1LT K5331M  
2½LT K5331L  
Dgr H:225-302-311+331-314-351-  
360D-370-372  
P:201-210-280-304+340-  
305+351+338-308+313



Use in conjunction with Karl Fischer reagent VC? (ROMIL K510?).  
Protect from atmospheric moisture.  
Application: Solvent (xylene-based) for volumetric KF titration of crude oils

### Karl Fischer solvent VS Hi-Dry KF

K6200

1LT K6200M  
2½LT K6200L  
Dgr H:225-301+311+331-314-360D-  
370  
P:201-210-280-302+352-309+310-  
403+235



Use in conjunction with Karl Fischer titrant VT? (ROMIL K621?).  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Solvent component (methanol-based) of pyridine-free 2-part system for volumetric KF titration (general purpose)

### Karl Fischer solvent VSA1.67 Hi-Dry KF

acid bu er

K6717

1LT K6717M  
Dgr H:225-301+311+331-314-360D-  
370  
P:201-210-280f-302+352-309+310-  
403+235



d 0.79  
Bu er capacity 1.67 mmol acid/ml  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Bu er solvent for volumetric KF titration

### Karl Fischer solvent VSA5 Hi-Dry KF

acid bu er

K6735

500ml K6735P  
1LT K6735M  
Dgr H:225-301+311+331-314-360D-  
370  
P:201-210-280-302+352-309+310-  
403+235



d 0.79  
Bu er capacity 5 mmol acid/ml  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Bu er solvent for volumetric KF titration

## ROMIL Hi-Dry® KF Karl Fischer Reagents and Calibrants Specifications

pyridine-free reagents for water determination

### Karl Fischer solvent VSO Hi-Dry KF

for oils

K6524

1LT K6524M  
Dgr H:225-301+311+331-314-360D-370  
P:201-210-280-302+352-309+310-403+235

Use in conjunction with Karl Fischer titrant VT? (ROMIL K621?).  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Solvent component of pyridine-free 2-part system for volumetric KF titration of oils



### Karl Fischer solvent VSOF Hi-Dry KF

for oils and fats

K6586

1LT K6586M  
2½LT K6586L  
Dgr H:226-302+312-331-314-351-360D-370-372  
P:201-210-261v-280F-304+340-305+351+338-308+313

Use in conjunction with Karl Fischer titrant VT? (ROMIL K621?).  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Solvent component (chloroform-based) of pyridine-free 2-part system for volumetric KF titration of oils and fats



### Karl Fischer titrant VT2 Hi-Dry KF

K6212

500ml K6212P  
1LT K6212M  
2½LT K6212L  
Dgr H:225-301+311+331-370-401  
P:210-280F-302+352-309+310-403+235

Nominal titre 1.00 ml = 2.00 mg ±1% H<sub>2</sub>O @ 20°C □ Use in conjunction with Karl Fischer solvent VS (ROMIL K6200).  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Titrant component of pyridine-free 2-part system for volumetric KF titration



### Karl Fischer titrant VT5 Hi-Dry KF

K6215

500ml K6215P  
1LT K6215M  
2½LT K6215L  
Dgr H:225-301+311+331-370-401  
P:210-280F-302+352-309+310-403+235

Nominal titre 1.00 ml = 5.00 mg ±0.5% H<sub>2</sub>O @ 20°C □ Use in conjunction with Karl Fischer solvent VS (ROMIL K6200).  
Protect from atmospheric moisture.  
Replace cap tightly immediately after use.  
Application: Titrant component of pyridine-free 2-part system for volumetric KF titration



### Karl Fischer calibrant BX5 Hi-Dry KF

H<sub>2</sub>O 5 mg/ml (water in m-xylene/butan-1-ol)

K7473

100ml K7473S  
500ml K7473P  
1LT K7473M  
Dgr H:226-302+312+332-315-318-335-336  
P:210-280-302+352-304+340-305+351+338

H<sub>2</sub>O 5.00 mg/ml (5000 ppm) ±0.02 mg/ml  
Matrix: m-Xylene/Butan-1-ol  
Protect from atmospheric moisture.  
Application: Reference material for KF titration (non-hygroscopic matrix)  
*SI-traceable through NIST certified reference material*



### Karl Fischer calibrant M5 Hi-Dry KF

H<sub>2</sub>O 5 mg/ml (water in methanol)

K7582

100ml K7582S  
500ml K7582P  
1LT K7582M  
Dgr H:225-301+311+331-370  
P:210-280F-302+352-309+310-403+235

d 0.79  
H<sub>2</sub>O 5 mg/ml (5000 ppm) ± 0.02 mg/ml  
Matrix: Methanol  
Protect from atmospheric moisture.  
Application: Reference material for KF titration  
*SI-traceable through NIST certified reference material*  
Supplied with Hi-Dry septum seal cap.



### Karl Fischer calibrant PC Hi-Dry KF

H<sub>2</sub>O 5.55% (potassium citrate)

K7755

10g K7755X

K3C6H5O7.H<sub>2</sub>O MW 324.41 CAS [6100-05-6]  
H<sub>2</sub>O 5.55% ±0.05%  
Dry at not greater than 105°C for 1 hour before use.  
Cool in desiccator.  
Application: Reference material for KF titration with oven

### Karl Fischer calibrant ST Hi-Dry KF

H<sub>2</sub>O 15.66% (sodium tartrate)

K7641

100g K7641S

Na<sub>2</sub>C<sub>4</sub>H<sub>4</sub>O<sub>6</sub>.2H<sub>2</sub>O MW 230.08 d 1.82 CAS [6106-24-7]  
H<sub>2</sub>O 15.66% ±0.05%  
Dry at not greater than 105°C for 1 hour before use.  
Cool in desiccator.  
Application: Reference material for KF titration

## ROMIL Hi-Dry® KF Karl Fischer Reagents and Calibrants Specifications

pyridine-free reagents for water determination

### Karl Fischer calibrant WS1 Hi-Dry KF

H<sub>2</sub>O 1 mg/g

K7129

100ml K7129S d 1.00  
 Wng H:226 Nominal concentration H<sub>2</sub>O 1.00 mg/g ±1% @ 20°C  
 P:210-262 Matrix: Anisole/Propylene Carbonate  
 Protect from atmospheric moisture.  
 Application: Reference material for KF titration  
 SI-traceable through NIST certified reference material  
 Supplied with Hi-Dry septum seal cap.



### Karl Fischer calibrant WS1 Hi-Dry KF

H<sub>2</sub>O 1 mg/g ampoule pack

K7142

10x5ml K7142Q d 1.00  
 Wng H:226 Nominal concentration H<sub>2</sub>O 1.00 mg/g ±2% @ 20°C  
 P:210-262 Matrix: Anisole/Propylene Carbonate  
 Application: Reference material for KF titration  
 SI-traceable through NIST certified reference material  
 Supplied in pack containing 10x5ml sealed glass ampoules.



### Karl Fischer calibrant WS10 Hi-Dry KF

H<sub>2</sub>O 10 mg/g

K7210

100ml K7210S d 1.00  
 Dgr H:226-315-318-335-336 Nominal concentration H<sub>2</sub>O 10.0 mg/g ±1% @ 20°C  
 P:210-280f-302+352-304+340- Matrix: Propylene Carbonate/p-Xylene/Butan-1-ol  
 305+351+338 Protect from atmospheric moisture.  
 Application: Reference material for KF titration  
 SI-traceable through NIST certified reference material  
 Supplied with Hi-Dry septum seal cap.



### Karl Fischer calibrant WS10 Hi-Dry KF

H<sub>2</sub>O 10 mg/g ampoule pack

K7264

10x10ml K7264Q d 1.00  
 Dgr H:226-315-318-335-336 Nominal concentration H<sub>2</sub>O 10.0 mg/g ±1% @ 20°C  
 P:210-280f-302+352-304+340- Matrix: Propylene Carbonate/p-Xylene/Butan-1-ol  
 305+351+338 Application: Reference material for KF titration  
 SI-traceable through NIST certified reference material  
 Supplied in pack containing 10 x 10ml sealed glass ampoules.



### Karl Fischer calibrant X0.1 Hi-Dry KF

H<sub>2</sub>O 0.1 mg/g (water in m-xylene)

K7362

100ml K7362S d 0.86  
 Wng H:226-312+332- H<sub>2</sub>O 0.100 mg/g ±2% @ 20°C  
 315 P:210-302+352- Matrix: m-Xylene  
 304+340 Protect from atmospheric moisture.  
 Application: Reference material for KF titration  
 SI-traceable through NIST certified reference material  
 Supplied with Hi-Dry septum seal cap.



### Karl Fischer calibrant X0.1 Hi-Dry KF

H<sub>2</sub>O 0.1 mg/g (water in m-xylene) ampoule pack

K7381

10x5ml K7381Q d 0.86  
 Wng H:226-312+332- H<sub>2</sub>O 0.100 mg/g ±5% @ 20°C  
 315 P:210-302+352- Matrix: m-Xylene  
 304+340 Protect from atmospheric moisture.  
 Application: Reference material for KF titration  
 SI-traceable through NIST certified reference material  
 Supplied in pack containing 10 x 5ml sealed glass ampoules.



Methyl Alcohol (see Methanol)

### Methanol Hi-Dry KF

K8408

1LT K8408M (Methyl Alcohol)  
 2½LT K8408L CH<sub>3</sub>OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
 Dgr H:225-301+311+331-370 Water <0.0100%  
 P:210-280f-302+352-309+310- Protect from atmospheric moisture.  
 403+235 Replace cap tightly immediately after use.  
 Application: General purpose solvent for KF sample dissolution



## ROMIL Hi-Dry® KF Karl Fischer Reagents and Calibrants Specifications

pyridine-free reagents for water determination

### Molecular Sieve 3A Hi-Dry KF

pellets 1.6mm

[K8678](#)

500g K8678P	d 0.64 CAS [1318-02-1]	□
1KG	Crystalline potassium aluminosilicate	
K8678M	Pore diameter about 3 Angstroms	
P:260d	Pellet size about 1.6mm (1/16 inch)	
	Protect from atmospheric moisture.	
	Application: Packing material for drying tubes in KF titration	

### Salicylic Acid Hi-Dry KF

[K8715](#)

500g K8715P	C <sub>6</sub> H <sub>4</sub> (OH)COOH MW138.12 CAS[69-72-7]	□
Dgr H:302-318	Water <0.2%	
P:280c-305+351+338-313	Protect from atmospheric moisture. Replace cap tightly immediately after use.	
	Application: Buffer reagent for KF titration	



### Xylene Hi-Dry KF

[K8982](#)

1LT K8982M	C <sub>6</sub> H <sub>4</sub> (CH <sub>3</sub> ) <sub>2</sub> MW106.17BP 138-142°C d 0.86 CAS [1330-20-7]	□
Wng H:226-312+332-315	Water <0.0050%	
P:210-302+352-304+340	Comprises 3 isomers and ethylbenzene	
	Application: Solvent for KF sample dissolution	



## ROMIL-SpR™ Super Purity Reagents Specifications


featuring ion-pair & buffer reagents, additives for LC-MS

**Ammonium Hydroxide solution** (see Ammonia solution)

**Ammonium Phosphate monobasic** (see Ammonium di-Hydrogen Phosphate)


### Acetic Acid SpR

HA016

100ml HA016S (Acetic Acid glacial)   
 500ml HA016P CH<sub>3</sub>COOH MW 60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7]  
 1LT HA016M Assay >99.8% Water <0.1% Residue <0.0001%  
 Dgr H:226-314 UV: 252nm >10%; 260nm >50%; 270nm >80%; 280nm >95%; 300-400nm >99%  
 P:280c-301+330+331-305+351+338- Fluorescence (0.1% aqueous, as quinine): 254nm <1 ppb; 365nm <1 ppb  
 307+310 Suitability for HPLC passes test  
 Suitability for LC-MS passes test  
 Trace ionic impurities (0.1% aqueous):  
 Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
 Al, Ca <25 ppb each  
 Na <50 ppb  
 Application: HPLC as buffer reagent, LC-MS as additive


### Ammonia solution SpR

HB059

50ml HB059T (Ammonium Hydroxide solution)   
 100ml HB059S NH<sub>3</sub> MW 17.03 d 0.92 CAS [1336-21-6]  
 Dgr H:314-335 Assay 20-22% Residue <0.0002%  
 P:280c-301+330+331-304+340- UV (0.1% aqueous): 225nm >20%; 235nm >50%; 240nm >80%; 250nm >95%;  
 305+351+338-309+310 260-400nm >99%  
 Fluorescence (0.1% aqueous, as quinine): 254nm <1 ppb; 365nm <1 ppb  
 Suitability for HPLC passes test  
 Suitability for LC-MS passes test  
 Trace ionic impurities (0.1% aqueous):  
 Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
 Al, Ca <25 ppb each  
 Na <50 ppb  
 Application: HPLC as buffer reagent, LC-MS as additive  
 Elemental impurities specified at time of manufacture.


### Ammonium Acetate SpR

HR079

100g HR079S CH<sub>3</sub>COONH<sub>4</sub> MW 77.08 CAS [631-61-8]   
 500g HR079P Assay >99%\*  
 UV (0.1% aqueous): 210nm >10%; 220nm >50%; 230nm >80%; 235nm >95%; 245-400nm >99%  
 \*on anhydrous substance  
 Fluorescence (0.1% aqueous, as quinine): 254nm <1 ppb; 365nm <1 ppb  
 Suitability for HPLC passes test  
 Suitability for LC-MS passes test  
 Chloride <0.0005%  
 Sulphate <0.001%  
 Nitrate <0.001%  
 Trace ionic impurities (0.1% aqueous):  
 Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
 Al, Ca <25 ppb each  
 Na <50 ppb  
 Deliquescent, hygroscopic. Replace cap promptly after use.  
 Application: HPLC as buffer reagent, LC-MS as additive


### Ammonium Carbonate SpR

HR143

100g HR143S NH<sub>4</sub>HCO<sub>3</sub>+NH<sub>2</sub>COONH<sub>4</sub> MW 157.13 d 0.90 CAS [8000-73-5]   
 500g HR143P UV: 230nm >10%; 235nm >50%; 240nm >80%; 250nm >95%; 310-400nm >99%  
 Dgr H:302-315-318 Mixture of ammonium hydrogen carbonate and ammonium carbamate in  
 P:280e-302+352-305+351+338-313 approximately equimolar proportions.  
 UV measured as 1M solution in water  
 Decomposes on exposure to air. Replace cap promptly after use. Application:  
 HPLC as buffer reagent

### Ammonium Formate SpR

HR305

100g HR305S HCOONH<sub>4</sub> MW 63.06 d 1.26 CAS [540-69-2]  
 500g HR305P Assay >98%  
 Wng H:315-319-335 UV (0.1% aqueous): 215nm >10%; 225nm >50%; 230nm >80%; 240nm >95%;  
 P:261d- 245-400nm >99%  
 305+351+338   
 Fluorescence (0.1% aqueous, as quinine): 254nm <1 ppb; 365nm <1 ppb  
 Suitability for HPLC passes test  
 Suitability for LC-MS passes test  
 Chloride <0.0005%  
 Sulphate <0.005%  
 Trace ionic impurities (0.1% aqueous):  
 Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
 Al, Ca <25 ppb each  
 Na <50 ppb  
 Deliquescent, hygroscopic. Replace cap promptly after use.  
 Application: HPLC as buffer reagent, LC-MS as additive



## ROMIL-SpR™ Super Purity Reagents Specifications

featuring ion-pair & buffer reagents, additives for LC-MS

### Ammonium di-Hydrogen Phosphate SpR

HR192

100g HR192S  
500g HR192P

**(Ammonium Phosphate monobasic)**  
NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> MW 115.03 CAS [7722-76-1]  
Assay >99%  
UV: 200nm >10%; 205nm >50%; 250nm >80%; 310nm >95%; 340-400nm >99%  
UV measured as 1M solution in water  
Application: HPLC as buffer reagent

### Butane-1-sulphonic Acid sodium salt SpR

IP204

5g IP204X  
25g IP204V  
100g IP204S  
1KG IP204M

**(Sodium n-Butyl-1-sulphonate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>SO<sub>3</sub>.Na MW 160.17 CAS [2386-54-1]  
Assay >98%  
UV: 200nm >96%  
UV measured as 0.005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds

### Cetyltrimethylammonium Bromide (see Hexadecyltrimethylammonium Bromide)

### Decane-1-sulphonic Acid sodium salt SpR


IP210

25g IP210V  
100g IP210S  
1KG  
IP210M

**(Sodium n-Decyl-1-sulphonate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>9</sub>SO<sub>3</sub>.Na MW 244.33 CAS [13419-61-9]  
Assay >98%  
UV: 200nm >96%  
UV measured as 0.005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds

### Diuroacetic Acid SpR



HA744

50ml HA744T  
100ml HA744S  
Dgr H:314  
P:280C-301+330+331-305+351+338-310  


**CHF<sub>2</sub>COOH** MW 96.03 FP -1°C BP 133°C d 1.53 CAS [381-73-7]  
Assay >97.5% Water <0.15% Residue <0.0001%  
UV (0.1% aqueous): 215nm >10%; 230nm >50%; 235nm >80%; 245nm >95%; 255-400nm >99%  
Fluorescence (0.1% aqueous, as quinine): 254nm <1 ppb; 365nm <1 ppb  
Suitability for HPLC passes test  
Suitability for LC-MS passes test  
Trace ionic impurities (0.1% aqueous):  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Store tightly closed in an upright position.  
Application: HPLC as ion-pair reagent, LC-MS as additive

### Diocetyl Sulphosuccinate sodium salt SpR

IP226

25g IP226V  
100g IP226S  
1KG  
Dgr IP:226M5-318  
P:280-305+351+338  


**(Sodium Diocetyl sulphosuccinate)**  
C<sub>20</sub>H<sub>37</sub>O<sub>7</sub>S.Na MW 444.56 CAS [577-11-7]  
Assay >98%  
UV: 254nm >95%  
UV measured as 0.005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds

### Dodecane-1-sulphonic Acid sodium salt SpR



IP212

25g IP212V  
100g IP212S  
1KG  
IP212M

**(Sodium n-Dodecyl-1-sulphonate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>SO<sub>3</sub>.Na MW 272.38 CAS [2386-53-0]  
Assay >98%  
UV: 200nm >90%  
UV measured as 0.0005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds

### Dodecyltrimethylammonium Bromide SpR

IP412

25g IP412V  
Dgr H:301-315-319-335-410  
P:273-302+352-305+351+338-309+310  


CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>N(CH<sub>3</sub>)<sub>3</sub>.Br MW 308.35 CAS [1119-94-4]  
Assay >98%  
UV: 254nm >90%  
UV measured as 0.1M solution in water  
Application: HPLC as ion-pair reagent for acidic compounds

## ROMIL-SpR™ Super Purity Reagents Specifications

featuring ion-pair & buffer reagents, additives for LC-MS

### Formic Acid SpR

HA353

100ml HA355S  
500ml HA353P  
1LT HA353M  
Dgr H:226-302-314-331-EUH071  
P:210-280c-301+330+331-305+351+338-310



HCOOH MW 46.03 FP 8.3°C BP 100.6°C d 1.22 CAS [64-18-6] □  
Assay >98% Residue <0.0001%  
UV (0.1% aqueous): 225nm <20%; 235nm <50%; 240nm <80%; 250nm <95%; 260-400nm <99%  
Fluorescence (0.1% aqueous, as quinine): 254nm <1 ppb; 365nm <1 ppb  
Suitability for HPLC passes test  
Suitability for LC-MS passes test  
Trace ionic impurities (0.1% aqueous):  
Ag, Cu, Fe, K, Mg, Mn, Ni, Pb, Zn <10 ppb each  
Al, Ca <25 ppb each  
Na <50 ppb  
Application: HPLC as bu er reagent, LC-MS as additive, Molecular Biology  
*Concentrated Formic Acid slowly decomposes to carbonmonoxideand waterand the pressure built up can cause an explosion of the sealed glass bottle.*  
*As a safety measure we t larger glass bottles with a closure featuring a venting valve. Bottles with a non-vented closure should have pressure released regularly.*

### Hepta uorobutyric Acid SpR

IP235

25ml IP235V  
Dgr H:314  
P:280c-301+330+331-305+351+338-309+310



**(Per uorobutyric Acid)**  
CF<sub>3</sub>(CF<sub>2</sub>)<sub>2</sub>COOH MW 214.04 CAS [375-22-4]  
Assay >99%  
UV: 230nm >70%; 240nm >90%; 254nm >99%  
UV measured as 0.01M solution in water  
Application: HPLC as ion-pair reagent for peptide separations, HPLC-MS as volatileion-pair reagent forbasic compounds

### Heptane-1-sulphonic Acid sodium salt SpR

IP207

5g IP207X  
25g IP207V  
100g IP207S  
1KG IP207M

**(Sodium n-Heptyl-1-sulphonate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>6</sub>SO<sub>3</sub>.NaMW202.27 CAS [22767-50-6]  
Assay >98%  
UV: 200nm >96%  
UV measured as 0.005M solution in water  
Application: HPLC asion-pairreagentforbasiccompounds

### Hexadecyltrimethylammonium Bromide

#### SpR

IP416

25g IP416V  
Wng H:302-315-319-335-410  
P:273-302+352-305+351+338



**(Cetyltrimethylammonium Bromide)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>15</sub>N(CH<sub>3</sub>)<sub>3</sub>.BrMW364.64 CAS [57-09-0]  
Assay >98%  
UV: 254nm >90%  
UV measured as 0.1M solution in water  
Application: HPLC as ion-pairreagentforacidiccompounds

### Hexane-1-sulphonic Acid sodium salt SpR

IP206

5g IP206X  
25g IP206V  
100g IP206S  
1KG IP206M

**(Sodium n-Hexyl-1-sulphonate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>SO<sub>3</sub>.NaMW188.22 CAS [2832-45-3]  
Assay >98%  
UV: 200nm >96%  
UV measured as 0.005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds

Orthophosphoric Acid (see Phosphoric Acid 85%)

### Octane-1-sulphonic Acid sodium salt SpR

IP208

5g IP208X  
25g IP208V  
100g IP208S  
1KG IP208M

**(Sodium n-Octyl-1-sulphonate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>7</sub>SO<sub>3</sub>.NaMW216.28 CAS [5324-84-5]  
Assay >98%  
UV: 200nm >96%  
UV measured as 0.005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds

Per uorobutyric Acid (see Hepta uorobutyric Acid)

Potassium Phosphate dibasic (see di-Potassium Hydrogen Phosphate)

Potassium Phosphate monobasic (see Potassium di-Hydrogen Phosphate)

## ROMIL-SpR™ Super Purity Reagents Specifications

featuring ion-pair & buffer reagents, additives for LC-MS

### Pentane-1-sulphonic Acid sodium salt SpR IP205

5g IP205X  
25g IP205V  
100g IP205S  
1KG IP205M

**(Sodium n-Pentyl-1-sulphonate)**  
 $\text{CH}_3(\text{CH}_2)_4\text{SO}_3\text{Na}$  MW 174.21 CAS [22767-49-3]  
 Assay >98%  
 UV: 200nm >96%  
 UV measured as 0.005M solution in water  
 Application: HPLC as ion-pair reagent for basic compounds

### Perchloric Acid 60% SpR HA622

100ml HA622S  
Dgr H:271-290-314  
P:210-221-280c-301+330+331-305+351+338-309+310

**HClO<sub>4</sub>** MW 100.46 d 1.54 CAS [7601-90-3]  
 Assay ca. 60%  
 UV: 254nm >98%  
 UV measured as 1M solution in water  
 Application: HPLC as buffer reagent



### Phosphoric Acid 85% SpR HA614

100ml HA614S  
500ml HA614P  
1LT HA614M  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-308+310

**(Orthophosphoric Acid)**  
 $\text{H}_3\text{PO}_4$  MW 98.00 d 1.70 CAS [7664-38-2]  
 Assay ca. 85%  
 UV (0.1% aqueous): 254nm >99%  
 Suitability for HPLC passes test  
 Application: HPLC as additive



### Potassium Bromide SpR HR706

for infra-red spectroscopy

100g HR706S

**KBr** MW 119.00 d 1.30 CAS [7758-02-3]  
 IR Spectroscopy passes test  
 Store in desiccator.  
 Application: IR Spectroscopy as support substance in fabrication of pressed sample discs

### Potassium di-Hydrogen Phosphate SpR HR739

100g HR739S  
500g HR739P

**(Potassium Phosphate monobasic)**  
 $\text{KH}_2\text{PO}_4$  MW 136.08 CAS [7778-77-0]  
 Assay >99.0%  
 UV: 200nm >10%; 205nm >50%; 210nm >80%; 305nm >95%; 320-400nm >98%  
 UV measured as 1M solution in water  
 Hygroscopic. Replace cap promptly after use.  
 Application: HPLC as buffer reagent

### di-Potassium Hydrogen Phosphate 3H<sub>2</sub>O SpR HR471

100g HR471S  
500g HR471P

**(Potassium Phosphate dibasic)**  
 $\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$  MW 228.22 CAS [16788-57-1]  
 Assay >99%  
 UV: 215nm >10%; 220nm >50%; 230nm >80%; 240nm >95%; 320-400nm >99%  
 UV measured as 1M solution in water  
 Adsorbs moisture from air. Replace cap promptly after use.  
 Application: HPLC as buffer reagent

**Sodium n-Butyl-1-sulphonate** (see [Butane-1-sulphonic Acid sodium salt](#))

**Sodium n-Decyl-1-sulphonate** (see [Decane-1-sulphonic Acid sodium salt](#))

**Sodium Dioctylsulphosuccinate** (see [Dioctyl Sulphosuccinate sodium salt](#))

**Sodium n-Dodecyl-1-sulphonate** (see [Dodecane-1-sulphonic Acid sodium salt](#))

**Sodium n-Heptyl-1-sulphonate** (see [Heptane-1-sulphonic Acid sodium salt](#))

**Sodium n-Hexyl-1-sulphonate** (see [Hexane-1-sulphonic Acid sodium salt](#))

**Sodium Lauryl Sulphate** (see [Sodium Dodecyl Sulphate](#))

**Sodium n-Octyl-1-sulphonate** (see [Octane-1-sulphonic Acid sodium salt](#))

**Sodium n-Pentyl-1-sulphonate** (see [Pentane-1-sulphonic Acid sodium salt](#))

**Sodium Propyl-1-sulphonate** (see [Propane-1-sulphonic Acid sodium salt](#))

## ROMIL-SpR™ Super Purity Reagents Specifications

featuring ion-pair & buffer reagents, additives for LC-MS

### Sodium Dodecyl Sulphate SpR

IP229

25g IP229V  
100g IP229S  
1KG  
Dgr H:228-302-319-308-412  
P:210h-273-280e-302+352-305+351+338-313

**(Sodium Lauryl Sulphate)**  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na MW 288.38 CAS [151-21-3]  
Assay >99%  
UV: 254nm >99%  
UV measured as 0.005M solution in water  
Application: HPLC as ion-pair reagent for basic compounds



### Sodium Perchlorate 1H<sub>2</sub>O SpR

HR343

100g HR343S  
500g HR343P  
Dgr H:271-302  
P:210-221

NaClO<sub>4</sub>.H<sub>2</sub>O MW 140.46 d 2.02 CAS [7791-07-3]  
Assay >98%  
UV: 210nm >80%; 220nm >90%; 230nm >95%; 240-400nm >98%  
UV measured as 1M solution in water  
Hygroscopic. Replace cap promptly after use.  
Application: HPLC as bu er reagent



Tetramethylene Sulphone (see Sulpholane)

Trichloromethane (see Chloroform)

### Tetrachloroethylene Hi-Dry

D4702

100ml D4702S  
500ml D4702P  
1LT D4702M  
2½LT D4702L  
Wng H:315-317-319-336-351-411  
P:273-281-302+352-305+351+338-308+313

**(Perchloroethylene)**  
CCl<sub>2</sub>CCl<sub>2</sub> MW 165.83 BP 121.1°C d 1.62 CAS [127-18-4]  
Assay >99.9% Water <0.0020% Residue <0.0001%  
Unstabilised



### Tetrahydrofuran Hi-Dry

D4718

100ml D4718S  
500ml D4718P  
1LT D4718M  
2½LT D4718L  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub> MW 72.11 BP 66°C d 0.89 CAS [109-99-9]  
Assay >99.9% Water <0.0025% Residue <0.0001%  
Unstabilised



### Tetrahydrofuran Hi-Dry

over molecular sieve

F8717

100ml F8717S  
500ml F8717P  
1LT F8717M  
2½LT F8717L  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66°C d 0.89 CAS [109-99-9]  
Assay >99.9% Water <0.0025%  
Unstabilised  
Contains molecular sieve.



### Tetrahydrofuran Hi-Dry

stabilised with BHT

D4719

100ml D4719S  
500ml D4719P  
1LT D4719M  
2½LT D4719L  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66°C d 0.89 CAS [109-99-9]  
Assay >99.9%\* Water <0.0025% Residue <0.0001%\*  
\*ex stabiliser  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm



### Toluene Hi-Dry

D4771

100ml D4771S  
500ml D4771P  
1LT D4771M  
2½LT D4771L  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235

C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub> MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]  
Assay >99.9% Water <0.0010% Residue <0.0001%



## ROMIL-Spr™ Super Purity Reagents Specifications

featuring ion-pair & buffer reagents, additives for LC-MS

### Toluene Hi-Dry

over molecular sieve

F8771

100ml F8771S  
500ml F8771P  
1LT F8771M  
2½LT F8771L  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235

C6H5CH3 MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]  
3] Assay >99.9% Water <0.0010%  
Contains molecular sieve.



### Triethylamine Hi-Dry

(CH<sub>3</sub>CH<sub>2</sub>)<sub>3</sub>N MW101.19BP88.9°Cd 0.73 CAS [121-44-8]  
Assay >99.8% Water <0.0050% Residue <0.0005%

D4763

100ml D4763S  
500ml D4763P  
1LT D4763M  
Dgr H:225-302+312+332-314-335  
P:210-280-301+330+331-302+352-304+340-305+351+338-309+310-403+235



### Triethylamine Hi-Dry

over molecular sieve

F8763

100ml F8763S  
500ml F8763P  
1LT F8763M  
Dgr H:225-302+312+332-314-335  
P:210-280-301+330+331-302+352-304+340-305+351+338-309+310-403+235

(CH<sub>3</sub>CH<sub>2</sub>)<sub>3</sub>N MW 101.19 BP 88.9°C d 0.73 CAS [121-44-8]  
8] Assay >99.8% Water <0.0050%  
Contains molecular sieve.



### 2,2,4-Trimethylpentane Hi-Dry

(iso-Octane)

(CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1]  
1] Assay >99.75% Water <0.0005% Residue <0.0001%

D4901

100ml D4901S  
500ml D4901P  
1LT D4901M  
2½LT D4901L  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-331-302+352-304+340-403+235



### 2,2,4-Trimethylpentane Hi-Dry

over molecular sieve

F8901

100ml F8901S  
500ml F8901P  
1LT F8901M  
2½LT F8901L  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-331-302+352-304+340-403+235

(CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1]  
1] Assay >99.75% Water <0.0005%  
Contains molecular sieve.



## ROMIL-BiO™ BioPure Solvents Specifications

for molecular biology

**Ammonium Hydroxide solution (see Ammonia solution)**

**Ammonium Hydroxide solution (see Ammonia solution)**

**iso-Amyl Alcohol (see 3-Methylbutan-1-ol)**

### Acetonitrile 10 BiO

H051

1LT H051M  
2½LT H051L  
4LT H051KZ  
BULK H051B  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

**(Methyl Cyanide)**  
CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8] Assay >99.9% Water <0.001% Residue <0.0001% Application: Protein sequencing, DNA synthesis



### Acetonitrile 30 BiO

H053

1LT H053M  
2½LT H053L  
4LT H053KZ  
BULK H053B  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

**(Methyl Cyanide)**  
CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8] Assay >99.9% Water <0.003% Residue <0.0001% Application: Primary washing agent in DNA synthesis



### Acetonitrile 50 BiO

H055

1LT H055M  
2½LT H055L  
4LT H055KZ  
BULK H055B  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

**(Methyl Cyanide)**  
CH<sub>3</sub>CN MW 41.05 BP 81.6°C d 0.78 CAS [75-05-8] Assay >99.9% Water <0.005% Residue <0.0001%



### Activator reagent ETT 3.25% BiO

**5-ethylthiotetrazole/acetonitrile**

D820

1LT D820M  
2½LT D820L  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

**(5-Ethylthiotetrazole solution)**  
Water <0.005%  
Contains:  
5-Ethylthiotetrazole 32.5 g/LT (0.25M)  
Solvent: Acetonitrile  
Application: Oligonucleotide Synthesis



### Ammonia solution BiO

H060

100ml H060S  
Dgr H:290-314-335-400  
P:273-280-301+330+331-304+340-305+351+338-309+310

**(Ammonium Hydroxide solution)**  
NH<sub>3</sub> MW 17.03 d 0.88 CAS[1336-21-6]  
Assay ca. 34% Residue <0.002%  
Colour <10 Hazen (APHA)  
Application: Molecular Biology



### Capping reagent A BiO

**acetic anhydride/lutidine/thf**

D519

1LT D519M  
2½LT D519L  
Dgr H:225-315-318-351-335  
P:210-240-280-302+352-305+351+338-308+313-403+233

Contains:  
Acetic Anhydride  
Lutidine  
Tetrahydrofuran  
Application: Oligonucleotide Synthesis



### Capping reagent B BiO

**N-methylimidazole/thf 16:84**

D576

1LT D576M  
2½LT D576L  
Dgr H:225-314-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

Water <0.015%  
Contains:  
N-Methylimidazole 16% v/v  
Tetrahydrofuran 84% v/v  
Application: Oligonucleotide Synthesis



## ROMIL-BiO™ BioPure Solvents Specifications

for molecular biology

Deblock reagent (see Detritylation reagent 3%)

### Detritylation reagent 3% BiO

dca/dichloromethane

D1682

1LT D1682M  
2½LT D1682L  
Wng H:315-319-351-402  
P:273-281-302+352-305+351+338-308+313

**(Deblock reagent 3%)**  
Contains:  
Dichloroacetic Acid 30 g/LT  
Solvent: Dichloromethane  
Application: Oligonucleotide Synthesis



### Dichloromethane BiO

stabilised with amylene

H203

1LT H203M  
2½LT H203L  
4LT H203KZ  
Wng H:351  
P:281-308+313

**(Methylene Dichloride)**  
CH<sub>2</sub>Cl<sub>2</sub>MW84.93BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.002% Residue <0.0001%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm  
Application: Molecular Biology



### Dimethylformamide BiO

H251

2½LT H251L  
4LT H251KZ  
Dgr H:360D-226-312+332-319  
P:201-210-302+352-305+351+338-308+313

HCON(CH<sub>3</sub>)<sub>2</sub> MW73.09BP153.0°Cd0.95 CAS [68-12-2]  
Assay >99.9% Water <0.03%  
Amines as CH<sub>3</sub>NH<sub>2</sub> <0.001% (<10 ppm)  
Application: Molecular Biology



### Dimethylformamide BiO

with molecular sieve

H254

2½LT H254L  
4LT H254KZ  
Dgr H:360D-226-312+332-319  
P:201-210-302+352-305+351+338-308+313

HCON(CH<sub>3</sub>)<sub>2</sub> MW 73.09 BP 153.0°C d 0.95 CAS [68-12-2]  
Assay >99.9% Water <0.003%  
Application: Molecular Biology



### Di-iso-propylethylamine BiO

H240

100ml H240S  
500ml H240P  
Dgr H:225-301-314-412  
P:210-233-240-273-280-301+330+331-305+351+338-309+310-403+235

**(Ethyl-di-iso-propylamine)**  
C<sub>8</sub>H<sub>19</sub>NMW 129.25 BP 127°C d 0.76 CAS [7087-68-5]  
Assay >99.5% Water <0.05% Residue <0.0001%  
Application: Molecular Biology



Ethyl-di-iso-propylamine (see Di-iso-propylethylamine)

5-Ethylthiotetrazole solution (see Activator reagent ETT)

Methyl Cyanide (see Acetonitrile)

Methylene Dichloride (see Dichloromethane)

### 3-Methylbutan-1-ol BiO

H440

100ml H440S  
500ml H440P  
Wng H:226-332-335-EUH066  
P:210-304+340

**(iso-Amyl Alcohol, iso-Pentyl Alcohol)**  
(CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>CH<sub>2</sub>OH MW88.15BP 131.1°C d 0.81 CAS [123-51-3]  
Assay >99.8% Water <0.005% Residue <0.0001%  
Comprises single isomer  
Application: Molecular Biology



### N-Methyl-2-pyrrolidone BiO

H567

2½LT H567L  
4LT H567KZ  
Dgr H:360D-315-319-335  
P:201-302+352-305+351+338-308+313





CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CONCH<sub>3</sub> MW99.13BP202.0°Cd1.03 CAS [872-50-4]  
Assay >99.5% Water <0.01%  
Application: Molecular Biology





## ROMIL-BiO™ BioPure Solvents Specifications

for molecular biology

1LT D552M 2½LT D532L Dgr H:225-319-335-351-EUH019 P:210-240-305+351+338-308+313-403+233	<b>Oxidiser reagent BiO</b> <b>iodine 0.02M in water/pyridine/thf</b> <span style="float: right;">D532</span> <hr/> Contains: <span style="float: right;">☐</span> Iodine 0.02M Water Pyridine Tetrahydrofuran Application: Oligonucleotide Synthesis
	<b>iso-Pentyl Alcohol (see 3-Methylbutan-1-ol)</b>
100ml H613S 500ml H613P Dgr H:225-300-311+331-314 P:210-280-301+330+331-302+352-304+340-305+351+338-309+310-403+235	<b>Piperidine BiO</b> <span style="float: right;">H613</span> <hr/> C5H11NMW 85.15 FP -11.0°C BP 106.2°C d 0.86 CAS [110-89-4] Assay >99.5% Water <0.05% Residue <0.0001% Application: Molecular Biology
	<b>Piperidine 20% BiO</b> <b>dmf solution</b> <span style="float: right;">D501</span> <hr/> Contains: Piperidine 20% v/v Dimethylformamide 80% v/v Application: DNA/RNA Synthesis
500ml D501P 1LT D501M 2½LT D501L Dgr H:226-311+331-314-360D P:201-210-302+352-305+351+338-308+313	<b>Pyridine BiO</b> <span style="float: right;">H649</span> <hr/> C5H5N MW 79.10BP 115.3°C d 0.98 CAS [110-86-1] Assay >99.8% Water <0.01% Residue <0.0002% Application: Molecular Biology
	<b>Pyridine BiO</b> <b>with molecular sieve</b> <span style="float: right;">H651</span> <hr/> C5H5N MW 79.10 BP 115.3°C CAS [110-86-1] Assay >99.8% Water <0.005% Application: Molecular Biology
2½LT H651L Dgr H:225-302+312+332-315-319 P:210-302+352-304+340-305+351+338-403+235	

## ROMIL-PUROM™ High Purity Process Solvents Specifications

for industrial applications

Acetic Acid glacial (see Acetic Acid)

### Acetic Acid PUROM

P5014

25LT P5014G  
200LT P5014D  
Dgr H:226-314  
P:280c-301+330+331-305+351+338-307+310

(Acetic Acid glacial)  
CH<sub>3</sub>COOH MW 60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7] Assay >99.8% Water <0.1% Residue <0.0005%  
Application: High purity process solvent



### Acetone PUROM

P5031

25LT P5031G  
200LT P5031D  
Dgr H:225-319-336-EUH066  
P:210-233-305+351+338

(Propanone)  
(CH<sub>3</sub>)<sub>2</sub>COMW 58.08 BP 56.1°C d 0.79 CAS [67-64-1]  
Assay >99.9% Water <0.2% Residue <0.0005%  
Application: High purity process solvent



### Acetonitrile PUROM

P5048

25LT P5048G  
200LT P5048D  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

(Methyl Cyanide)  
CH<sub>3</sub>CNMW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.01% Residue <0.0005%  
Application: High purity process solvent



### Acetonitrile PUROM

for preparative HPLC

P5046

25LT P5046G  
200LT P5046D  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233

(Methyl Cyanide)  
CH<sub>3</sub>CNMW 41.05 BP 81.6°C d 0.78 CAS [75-05-8]  
Assay >99.9% Water <0.03% Residue <0.0005%  
UV: 230nm >70%; 280nm >95%  
Application: High purity process solvent for preparative HPLC



n-Butanol (see Butan-1-ol)

n-Butyl Alcohol (see Butan-1-ol)

tert-Butyl Methyl Ether (see Methyl tert-Butyl Ether)

### Butan-1-ol PUROM

P5083

25LT P5083G  
200LT P5083D  
Dgr H:226-302-315-318-335-336  
P:210-280F-302+352-304+340-305+351+338-313

(n-Butanol, n-Butyl Alcohol)  
CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>OH MW74.12 BP 117.7°C d 0.81 CAS [71-36-3]  
Assay >99.8% Water <0.05% Residue <0.0005%  
Application: High purity process solvent



### Chloroform PUROM

stabilised with amylene

P5140

200LT P5140D  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280F-304+340-305+351+338-308+313

(Trichloromethane)  
CHCl<sub>3</sub> MW119.38 BP 61.2°C d 1.48 CAS [67-66-3]  
Assay >99.9%\* Water <0.005% Residue <0.0005%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm  
Application: High purity process solvent



### Chloroform PUROM

stabilised with ethanol

P5135

200LT P5135D  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280F-304+340-305+351+338-308+313

(Trichloromethane)  
CHCl<sub>3</sub> MW119.38 BP 61.2°C d 1.48 CAS [67-66-3]  
Assay >99.9%\* Water <0.005% Residue <0.0005%  
\*ex stabiliser  
Stabiliser: Ethanol ca. 1% w/w  
Stabiliser should only be removed immediately before use by adsorption onto activated alumina.  
Application: High purity process solvent



## ROMIL-PUROM™ High Purity Process Solvents Specifications

for industrial applications

### Dichloromethane PUROM

stabilised with amylene

P5202

25LT P5202G  
200LT P5202D  
Wng H:351  
P:281-308+313



**(Methylene Dichloride)**  
CH2Cl2 MW 84.93 BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99.9%\* Water <0.01% Residue <0.0005%  
\*ex stabiliser  
Stabiliser: Amylene ca. 25 ppm  
Application: High purity process solvent

### Diethyl Ether PUROM

stabilised with BHT

P5220

25LT P5220G  
200LT P5220D  
Dgr H:224-302-336-EUH019-EUH066  
P:210-240-403+235



**(C2H5)2O** MW 74.12 BP 34.4°C d 0.71 CAS [60-29-7]  
Assay >99.9%\* Water <0.02% Residue <0.0005%\*  
\*ex stabiliser  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 5 ppm  
Application: High purity process solvent

### 1,2-Dimethoxyethane PUROM

P5261

25LT P5261G  
200LT P5261D  
Dgr H:225-360FD-332-EUH019  
P:201-210-308+313-403+235



**(Ethylene Glycol Dimethyl Ether)**  
CH3OCH2CH2OCH3 MW 90.12 FP -58°C BP 85°C d 0.87 CAS [110-71-4]  
Assay >99.9% Water <0.01% Residue <0.0005%  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Application: High purity process solvent

### Dimethylformamide PUROM

P5253

25LT P5253G  
200LT P5253D  
Dgr H:360D-226-312+332-319  
P:201-210-302+352-305+351+338-308+313



**HCON(CH3)2** MW 73.09 BP 153.0°C d 0.95 CAS [68-12-2]  
Assay >99.9% Water <0.03% Residue <0.0005%  
Application: High purity process solvent

### 1,4-Dioxan PUROM

P5297

25LT P5297G  
200LT P5297D  
Dgr H:225-350-319-335-EUH019-EUH066  
P:210-281-305+351+338-308+313



**C4H8O2** MW 88.11 FP 11.8°C BP 101.3°C d 1.03 CAS [123-91-1]  
Assay >99.9% Water <0.005% Residue <0.0005%  
Unstabilised  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Application: High purity process solvent

Ethyl Alcohol (see Ethanol)

Ethylene Glycol Dimethyl Ether (see 1,2-Dimethoxyethane)

### Ethanol absolute PUROM

P5314

25LT P5314G  
200LT P5314D  
Dgr H:225  
P:210-233-240-403+235



**(Ethyl Alcohol)**  
C2H5OH MW 46.07 BP 78.3°C d 0.79 CAS [64-17-5]  
Assay >99.8% Water <0.1% Residue <0.0005%  
Application: High purity process solvent

### Ethyl Acetate PUROM

P5346

25LT P5346G  
200LT P5346D  
Dgr H:225-319-336-EUH066  
P:210-233-240-305+351+338-403+235



**CH3COOC2H5** MW 88.11 BP 77.1°C d 0.90 CAS [141-78-6]  
Assay >99.9% Water <0.005% Residue <0.0005%  
Application: High purity process solvent

### n-Heptane 95% PUROM

P5367

25LT P5367G  
200LT P5367D  
Dgr H:225-304-315-336-410  
P:210-273-301+310-331-302+352-304+340-403+235



**CH3(CH2)5CH3** MW 100.21 BP 94-98°C d 0.68 CAS [142-82-5]  
Water <0.005% Residue <0.0005%  
Assay (n-isomer) >95%  
Assay (all isomers) >99.5%  
Application: High purity process solvent

## ROMIL-PUROM™ High Purity Process Solvents Specifications

for industrial applications

25LT P5366G 200LT P5366D Dgr H:225-304-315-336-410 P:210-273-301+310-331-302+352-304+340-403+235	<b>n-Heptane 99% PUROM</b> <span style="float:right">P5366</span> <hr/> CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CH <sub>3</sub> MW 100.21 BP98.4°C d0.68 CAS [142-82-5] <span style="float:right">☐</span> Assay >99% Water <0.005% Residue <0.0005% Application: High purity process solvent
	<b>Hexane fraction PUROM</b> <span style="float:right">P5390</span> <hr/> C <sub>6</sub> H <sub>14</sub> BP 65-70°C d 0.66 CAS [73513-42-5] <span style="float:right">☐</span> Water <0.005% Residue <0.0005% Comprises ca. 50% n-isomer, the remainder being predominantly other isomers of hexane. Application: High purity process solvent
	<b>iso-Hexane 95% PUROM</b> <span style="float:right">P5388</span> <hr/> C <sub>6</sub> H <sub>14</sub> MW86.18BP55-63°Cd0.65CAS[107-83-5] <span style="float:right">☐</span> Water <0.005% Residue <0.0005% n-Hexane <5% Application: High purity process solvent
	<b>n-Hexane 95% PUROM</b> <span style="float:right">P5389</span> <hr/> CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> CH <sub>3</sub> MW 86.18BP67-70°Cd 0.66 CAS [110-54-3] <span style="float:right">☐</span> Water <0.005% Residue <0.0005% Assay (n-isomer) >95% Assay (all isomers) >99.5% Application: High purity process solvent
	<b>Methyl Alcohol (see Methanol)</b> <b>Methyl Cyanide (see Acetonitrile)</b> <b>Methylene Dichloride (see Dichloromethane)</b> <b>4-Methylpentan-2-one (see Methyl iso-Butyl Ketone)</b>
25LT P5410G 200LT P5410D Dgr H:225-301+311+331-370 P:210-280f-302+352-309+310-403+235	<b>Methanol PUROM</b> <span style="float:right">P5410</span> <hr/> <b>(Methyl Alcohol)</b> <span style="float:right">☐</span> CH <sub>3</sub> OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1] Assay >99.9% Water <0.02% Residue <0.0005% Application: High purity process solvent
	<b>Methanol PUROM for preparative HPLC</b> <span style="float:right">P5408</span> <hr/> <b>(Methyl Alcohol)</b> <span style="float:right">☐</span> CH <sub>3</sub> OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1] Assay >99.8% Water <0.05% Residue <0.0005% UV: 225nm >50%; 255nm >95% Application: High purity process solvent for preparative HPLC
	<b>Methyl tert-Butyl Ether PUROM</b> <span style="float:right">P5447</span> <hr/> <b>(tert-Butyl Methyl Ether)</b> <span style="float:right">☐</span> CH <sub>3</sub> OC(CH <sub>3</sub> ) <sub>3</sub> MW 88.15 BP 55.4°C d 0.74 CAS [1634-04-4] Assay >99.7% Water <0.02% Residue <0.0005% Peroxides (at time of manufacture) <0.0001% (<1 ppm) Application: High purity process solvent
	<b>2-Methyltetrahydrofuran PUROM</b> <span style="float:right">P5536</span> <hr/> CH <sub>3</sub> C <sub>4</sub> H <sub>7</sub> OMW86.13 BP80°Cd0.86 CAS[96-47-9] Assay >99.8% Water <0.01% Residue <0.0005% Unstabilised Peroxides (at time of manufacture) <0.0001% (<1 ppm) Application: High purity process solvent
	


## ROMIL-PUROM™ High Purity Process Solvents Specifications

for industrial applications

### 2-Methyltetrahydrofuran PUROM

stabilised with BHT

P5537

25LT P5537G CH<sub>3</sub>C<sub>4</sub>H<sub>7</sub>O MW 86.13 BP 80°C d 0.86 CAS [96-47-9]   
 200LT P5537D Assay >99.8%\* Water <0.01% Residue <0.0005%\*  
 Dgr H:225-319-335-EUH019 \*ex stabiliser  
 P:210-233-240-305+351+338- Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
 403+235 Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm  
 Application: High purity process solvent



#### iso-Octane (see 2,2,4-Trimethylpentane)

Perchloroethylene (see Tetrachloroethylene)

Petroleum Distillate (see Petroleum Ether)

Petroleum Spirit (see Petroleum Ether)

n-Propanol (see Propan-1-ol)

iso-Propanol (see Propan-2-ol)


Propanone (see Acetone)

n-Propyl Alcohol (see Propan-1-ol)

iso-Propyl Alcohol (see Propan-2-ol)

### n-Pentane 95% PUROM


P5571

25LT P5571G CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub> MW 72.15BP35.5-37°Cd 0.63 CAS [109-66-0]   
 200LT P5571D Water <0.005% Residue <0.0005%  
 Dgr H:225-304-336-411-EUH066 Assay (n-isomer) >95%  
 P:273-301+310-331-403+235 Assay (all isomers) >99.5%  
 Application: High purity process solvent



### Petroleum Ether 40-60°C PUROM


P5601

25LT P5601G (Petroleum Distillate, Petroleum Spirit)   
 200LT P5601D BP40-60°C d 0.64 CAS [8032-32-4]  
 Dgr H:225-304-336-411-EUH066 Water <0.005% Residue <0.0005%  
 P:210-233-243-273-280-301+310- Application: High purity process solvent  
 303+361+353-304+340-331-  
 403+235



### Petroleum Ether 60-80°C PUROM


P5602

25LT P5602G (Petroleum Distillate, Petroleum Spirit)   
 200LT P5602D BP60-80°C d 0.67  
 Dgr H:225-304-315-336-411 Water <0.005% Residue <0.0005%  
 P:210-243-273-280-301+310-331- Application: High purity process solvent  
 403+235



### Propan-1-ol PUROM


P5624

25LT P5624G (n-Propanol, n-Propyl Alcohol)   
 200LT P5624D CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>OH MW 60.10 BP 97.2°C d 0.80 CAS [71-23-  
 8] Assay >99.9% Water <0.05% Residue <0.0005%  
 Dgr H:225-318-336 Application: High purity process solvent  
 P:210-233-280F-305+351+338-313



### Propan-2-ol PUROM

P5625

25LT P5625G (iso-Propanol, iso-Propyl Alcohol)   
 200LT P5625D (CH<sub>3</sub>)<sub>2</sub>CHOH MW 60.10 BP 82.2°C d 0.78 CAS [67-63-0]  
 Dgr H:225-319-336 Assay >99.9% Water <0.02% Residue <0.0005%  
 P:210-233-305+351+338 Application: High purity process solvent



## ROMIL-PUROM™ High Purity Process Solvents Specifications

for industrial applications

Trichloromethane (see Chloroform)

### Tetrachloroethylene PUROM

P5702

25LT P5702G  
200LT P5702D  
Wng H:315-317-319-336-351-411  
P:273-281-302+352-305+351+338-308+313

**(Perchloroethylene)**  
CCl<sub>2</sub>CCl<sub>2</sub>MW 165.83 BP 121.1°C d 1.62 CAS [127-18-4]  
Assay >99.9% Water <0.005% Residue <0.0005%  
Unstabilised  
Application: High purity process solvent



### Tetrahydrofuran PUROM

P5718

25LT P5718G  
200LT P5718D  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>OMW 72.11BP 66.0°C d0.89 CAS [109-99-9]  
Assay >99.9% Water <0.005% Residue <0.0005%  
Unstabilised  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Application: High purity process solvent



### Tetrahydrofuran PUROM

stabilised with BHT

P5719

25LT P5719G  
200LT P5719D  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66.0°C d 0.89 CAS [109-99-9]  
Assay >99.9%\* Water <0.005% Residue <0.0005%\*  
\*ex stabiliser  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm  
Application: High purity process solvent



### Toluene PUROM

P5771

25LT P5771G  
200LT P5771D  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235

C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]  
Assay >99.9% Water <0.01% Residue <0.0005%  
Application: High purity process solvent



### 2,2,4-Trimethylpentane PUROM

P5901

25LT P5901G  
200LT P5901D  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-331-302+352-304+340-403+235

**(iso-Octane)**  
(CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1]  
Assay >99.75%Water <0.005% Residue<0.0005%  
Application:Highpurityprocess solvent



### Water PUROM

P5950

25LT P5950G  
200LT P5950D

H<sub>2</sub>OMW 18.02 FP 0.0°C BP 100.0°C CAS [7732-18-5]  
Residue <0.0001%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
Application: High purityprocess solvent

### Water PUROM

for preparative HPLC

P5948

25LT P5948G  
200LT P5948D

H<sub>2</sub>O MW 18.02 FP 0.0°C BP 100.0°C d 1.00 CAS [7732-18-5]  
Residue <0.0005%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
TOC (at time of manufacture) <50 ppb  
Application: High purity process solvent for preparative HPLC

### Xylene mixed isomers PUROM

P5982

25LT P5982G  
200LT P5982D  
Wng H:226-312+332-315  
P:210-302+352-304+340

C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW106.17BP138-142°Cd0.86CAS[1330-20-7]  
Water <0.01% Residue <0.0005%  
Comprises 3 isomers and ethylbenzene  
Assay (total C<sub>8</sub>H<sub>10</sub> isomers) >98.5%  
Ethylbenzene typically <3%  
Toluene typically <0.5%  
Methylethylbenzene typically <0.5%  
Application: High purity process solvent



## ROMIL-PUROM™ High Purity Process Solvents Specifications

for industrial applications

Trichloromethane (see Chloroform)

### Tetrachloroethylene PUROM

P5702

25LT P5702G  
200LT P5702D  
Wng H:315-317-319-336-351-411  
P:273-281-302+352-305+351+338-308+313

**(Perchloroethylene)**  
CCl<sub>2</sub>CCl<sub>2</sub>MW 165.83 BP 121.1°C d 1.62 CAS [127-18-4]  
Assay >99.9% Water <0.005% Residue <0.0005%  
Unstabilised  
Application: High purity process solvent



### Tetrahydrofuran PUROM

P5718

25LT P5718G  
200LT P5718D  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>OMW 72.11BP 66.0°C d0.89 CAS [109-99-9]  
Assay >99.9% Water <0.005% Residue <0.0005%  
Unstabilised  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Application: High purity process solvent



### Tetrahydrofuran PUROM

stabilised with BHT

P5719

25LT P5719G  
200LT P5719D  
Dgr H:225-319-335-351-EUH019  
P:210-240-305+351+338-308+313-403+233

CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>O MW 72.11 BP 66.0°C d 0.89 CAS [109-99-9]  
Assay >99.9%\* Water <0.005% Residue <0.0005%\*  
\*ex stabiliser  
Peroxides (at time of manufacture) <0.0001% (<1 ppm)  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm  
Application: High purity process solvent



### Toluene PUROM

P5771

25LT P5771G  
200LT P5771D  
Dgr H:225-304-315-336-361d-373  
P:210-240-301+310-331-302+352-403+235

C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>MW 92.14 BP 110.6°C d 0.87 CAS [108-88-3]  
Assay >99.9% Water <0.01% Residue <0.0005%  
Application: High purity process solvent



### 2,2,4-Trimethylpentane PUROM

P5901

25LT P5901G  
200LT P5901D  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-331-302+352-304+340-403+235

**(iso-Octane)**  
(CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1]  
Assay >99.75%Water <0.005% Residue<0.0005%  
Application:Highpurityprocess solvent



### Water PUROM

P5950

25LT P5950G  
200LT P5950D

H<sub>2</sub>OMW 18.02 FP 0.0°C BP 100.0°C CAS [7732-18-5]  
Residue <0.0001%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
Application: High purityprocess solvent

### Water PUROM

for preparative HPLC

P5948

25LT P5948G  
200LT P5948D

H<sub>2</sub>O MW 18.02 FP 0.0°C BP 100.0°C d 1.00 CAS [7732-18-5]  
Residue <0.0005%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
pH (at time of manufacture) 5.5-8.0 @ 25°C  
TOC (at time of manufacture) <50 ppb  
Application: High purity process solvent for preparative HPLC

### Xylene mixed isomers PUROM

P5982

25LT P5982G  
200LT P5982D  
Wng H:226-312+332-315 P:210-302+352-304+340

C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW106.17BP138-142°Cd0.86CAS[1330-20-7]  
Water <0.01% Residue <0.0005%  
Comprises 3 isomers and ethylbenzene  
Assay (total C<sub>8</sub>H<sub>10</sub> isomers) >98.5%  
Ethylbenzene typically <3%  
Toluene typically <0.5%  
Methylethylbenzene typically <0.5%  
Application: High purity process solvent





## ROMIL-SpA™ Super Purity Acids and Reagents Specifications

Acetic Acid glacial (see Acetic Acid)

Ammonium Hydroxide solution (see Ammonia solution)

### Acetic Acid SpA

H015

500ml H015P  
Dgr H:226-314  
P:280c-301+330+331-305+351+338-307+310



(Acetic Acid glacial)

CH<sub>3</sub>COOH MW60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7]

Assay >99%

Colour <10 Hazen (APHA)

Be, Bi, Ce, Co, Cs, Dy, Er, Eu, Ga, Gd, Hf, Ho, In, La, Li, Lu, Nd, Pb, Pr, Rb, Re, Sc, Sm, Tb, Th, Tl, Tm, U, Y, Yb, Zr <0.1 ppb each

As, Ba, Cd, Cu, Ge, Mg, Mn, Mo, Ni, Pt, Rh, Ru, Sb, Sn, Sr, Te, Ti, V, W <0.5 ppb each

Ag, Al, Ca, Cr, Fe, Hg, K, Na, Se, Zn <1 ppb each

SO<sub>4</sub> <0.5 ppm

PO<sub>4</sub> <1 ppm

Cl <1 ppm

Substances reducing dichromate passes test

Substances reducing permanganate passes test

Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis (for organic trace analysis see ROMIL Code H014 Acetic Acid SpS), Ion Chromatography, Voltammetry

*Elemental impurities specification at time of manufacture.*

*Typical values, elemental impurities at time of manufacture (ppb):*

Ag <0.1 Cu <0.2 K <0.5 Pt <0.1 Te <0.1 Al <0.5 Dy <0.1 La <0.1 Rb <0.1 Th <0.1 As <0.1 Er <0.1 Li <0.1 Re <0.1 Ti <0.1 Ba <0.1 Eu <0.1 Lu <0.1 Rh <0.1 Tl <0.1 Be <0.1 Fe <0.5 Mg <0.2 Ru <0.1 Tm <0.1 Bi <0.1 Ga <0.1 Mn <0.1 Sb <0.1 U <0.1 Ca <1 Gd <0.1 Mo <0.1 Sc <0.1 V <0.1 Cd <0.1 Ge <0.1 Na <1 Se <0.5 W <0.1 Ce <0.1 Hf <0.1 Nd <0.1 Sm <0.1 Y <0.1 Co <0.1 Hg <0.1 Ni <0.1 Sn <0.1 Yb <0.1 Cr <0.1 Ho <0.1 Pb <0.1 Sr <0.1 Zn <0.5 Cs <0.1 In <0.1 Pr <0.1 Tb <0.1 Zr <0.1

### Ammonia solution SpA

H058

500ml H058P  
Dgr H:314-335  
P:280c-301+330+331-304+340-305+351+338-309+310



(Ammonium Hydroxide solution)

NH<sub>3</sub> MW 17.03 d 0.92 CAS [1336-21-6]

Assay 20-22%

Colour <10 Hazen (APHA)

Ba, Be, Bi, Ce, Cs, Dy, Er, Eu, Ga, Gd, Ge, Ho, In, La, Li, Lu, Nb, Nd, Pb, Pr, Rb, Sc, Sm, Sr, Tb, Te, Th, Tl, Tm, U, W, Y, Yb, Zr <0.1 ppb each

Hg <0.2 ppb

Ag, Au, Cd, Co, Cr, Cu, Mn, Mo, Ni, Rh, Sb, Sn, Ti, V, Zn <0.5 ppb each

Al, As, Ca, Fe, K, Mg, Na, Se <1 ppb each

PO<sub>4</sub> <0.01 ppm

Cl <0.5 ppm

SO<sub>4</sub> <1 ppm

Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry

*Elemental impurities specification at time of manufacture.*

*Typical values, elemental impurities at time of manufacture (ppb):*

Ag <0.1 Cu <0.5 La <0.1 Pt <1 Th <0.1 Al <0.5 Dy <0.1 Li <0.1 Rb <0.1 Ti <0.1 As <0.1 Er <0.1 Lu <0.1 Re <1 Tl <0.1 Au <0.1 Eu <0.1 Mg <0.2 Rh <0.1 Tm <0.1 Ba <0.1 Fe <0.5 Mn <0.2 Ru <1 U <0.1 Be <0.1 Ga <0.1 Mo <0.1 Sb <0.1 V <0.1 Bi <0.1 Gd <0.1 Na <0.5 Sc <0.1 W <0.1 Ca <0.5 Ge <0.1 Nb <0.1 Se <0.1 Y <0.1 Cd <0.1 Hf <1 Nd <0.1 Sm <0.1 Yb <0.1 Ce <0.1 Hg <0.2 Ni <0.2 Sn <0.1 Zn <0.5 Co <0.1 Ho <0.1 Pb <0.1 Sr <0.1 Zr <0.1 Cr <0.1 In <0.1 Pd <1 Tb <0.1 Cs <0.1 K <0.2 Pr <0.1 Te <0.1

### Dimethylformamide SpA

low metals

H255

1LT H255M  
2½LT H255L  
Dgr H:360D-226-312+332-319  
P:201-210-302+352-305+351+338-308+313



HCON(CH<sub>3</sub>)<sub>2</sub> MW 73.09 BP 153.0°C d 0.95 CAS [68-12-2] Assay >99.9% Water <0.05% Residue <0.0001%

Group 1 & 2 elements typically <0.1-<10 ppb

Group 3 to 12 (transition) elements typically <0.1-<5 ppb

Group 13, 14, 15 elements typically <0.1-<5 ppb

Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis (for organic analysis and chromatography see ROMIL Code H253 Dimethylformamide SpS)

## ROMIL-SpA™ Super Purity Acids and Reagents Specifications

### Hydrochloric Acid SpA

H396

500ml H396P  
1LT H396M  
2½LT H396L  
Dgr H:290-314-335  
P:280c-301+330+331-305+351+338-309+310



HCl MW 36.46 d 1.18 CAS[7647-01-0] □ Assay 34-37%  
Colour <10 Hazen (APHA)  
Ba, Be, Bi, Cd, Ce, Co, Cs, Dy, Er, Eu, Ga, Gd, Hf, Ho, In, La, Li, Lu, Mn, Mo, Nb, Nd, Pb, Pr, Rb, Re, Rh, Ru, Sc, Sm, Sr, Tb, Te, Th, Tl, Tm, U, W, Y, Yb, Zr <0.1 ppb each  
As, Au, Cr, Cu, Hg, Mg, Ni, Sb, Sn, Ti, V <0.5 ppb each  
Ag, Al, B, Ca, Fe, K, Na, Se, Zn <1 ppb each  
Total P <0.01 ppm  
Total S <0.3 ppm  
Free Cl2 <0.5 ppm  
Br <10 ppm  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry  
*Elemental impurities specification at time of manufacture.*  
*Typical values, elemental impurities at time of manufacture (ppb):*  
Ag <0.1 Cs <0.1 La <0.1 Pt <0.5 Te <0.1 Al <0.5 Cu <0.1 Li <0.1 Rb <0.1 Th <0.1 As <0.1 Dy <0.1 Lu <0.1 Re <0.1 Ti <0.1 Au <0.1 Er <0.1 Mg <0.5 Rh <0.1 Tl <0.1 B <0.5 Eu <0.1 Mn <0.1 Ru <0.1 Tm <0.1 Ba <0.1 Fe <0.5 Mo <0.1 Sb <0.1 U <0.1 Be <0.1 Ga <0.1 Na <0.5 Sc <0.1 V <0.1 Bi <0.1 Gd <0.1 Nb <0.1 Se <0.1 W <0.1 Ca <0.5 Hf <0.1 Nd <0.1 Sm <0.1 Y <0.1 Cd <0.1 Hg <0.2 Ni <0.1 Sn <0.1 Yb <0.1 Ce <0.1 Ho <0.1 Pb <0.1 Sr <0.1 Zn <0.5 Co <0.1 In <0.1 Pd <0.5 Ta <0.5 Zr <0.1 Cr <0.1 K <0.1 Pr <0.1 Tb <0.1

### Hydrofluoric Acid SpA

H405

500ml H405P  
Dgr H:300+310+330-314  
P:280-301+330+331-302+352-304+340-305+351+338-310+R0P010



HF MW 20.01 d 1.16 CAS[7664-39-3] □ Assay 47-51%  
Colour <10 Hazen (APHA)  
Ba, Be, Bi, Cd, Ce, Co, Cs, Dy, Er, Eu, Ga, Gd, Ge, Hf, Ho, In, La, Li, Lu, Mn, Mo, Nb, Nd, Pb, Pr, Rb, Re, Rh, Ru, Sc, Sm, Sr, Tb, Te, Th, Tl, Tm, U, V, Y, Yb, Zr <0.1 ppb each  
Au, Pd, Pt, Sb <0.2 ppb each  
Ag, As, Cu, Ni, Sn, W <0.5 ppb each  
Al, B, Ca, Cr, Fe, Hg, K, Mg, Na, Se, Ti, Zn <1 ppb each  
Total P <0.05 ppm  
Total S <0.1 ppm  
Cl <4 ppm  
SiF6 <20 ppm  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry  
*Elemental impurities specification at time of manufacture.*  
*For treatment of HF burns, calcium gluconate gel (ROMIL Code PCG9V) is recommended. In order to provide emergency first aid, it should be kept wherever HF is handled or stored.*  
*Typical values, elemental impurities at time of manufacture (ppb):*  
Ag <0.1 Cs <0.1 K <0.2 Pr <0.1 Tb <0.1 Al <0.5 Cu <0.1 La <0.1 Pt <0.2 Te <0.1 As <0.1 Dy <0.1 Li <0.1 Rb <0.1 Th <0.1 Au <0.1 Er <0.1 Lu <0.1 Re <0.1 Ti <0.5 B <0.5 Eu <0.1 Mg <0.2 Rh <0.1 Tl <0.1 Ba <0.1 Fe <0.5 Mn <0.1 Ru <0.1 Tm <0.1 Be <0.1 Ga <0.1 Mo <0.1 Sb <0.1 U <0.1 Bi <0.1 Gd <0.1 Na <0.5 Sc <0.1 V <0.1 Ca <0.5 Ge <0.1 Nb <0.1 Se <0.1 W <0.5 Cd <0.1 Hf <0.1 Nd <0.1 Sm <0.1 Y <0.1 Ce <0.1 Hg <0.05 Ni <0.1 Sn <0.1 Yb <0.1 Co <0.1 Ho <0.1 Pb <0.1 Sr <0.1 Zn <0.1 Cr <0.1 In <0.1 Pd <0.2 Ta <0.5 Zr <0.1

### Hydrogen Peroxide SpA

H416

500ml H416P  
Dgr H:302-318  
P:280e-305+351+338-313



H2O2 MW 34.01d 1.10 CAS[7722-84-1]  
Assay ca. 30%  
Colour <10 Hazen (APHA)  
B, Be, Bi, Ce, Co, Cs, Dy, Er, Eu, Ga, Gd, Hf, Ho, Ir, La, Lu, Mn, Nd, Pd, Pt, Rb, Re, Rh, Ru, Sc, Sm, Sr, Tb, Te, Th, Tl, Tm, U, Y, Yb <0.5 ppb each  
Ba, Cu, Cr, Sb <1 ppb each  
Ag, As, Cd, Ge, In, Li, Mg, Mo, Ni, Pb, Se, Sn, V, Zr <5 ppb each  
Al, Ti, Zn <10 ppb each  
Au, Ca, Fe, Hg, K, Na, Nb, Ta, W <50 ppb each  
Cl, PO4 <0.5 ppm each  
SO4, NO3 <1 ppm each  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry  
*Elemental impurities specification at time of manufacture.*  
*As a safety measure we bottle with a closure featuring a venting valve.*  
*Typical values, elemental impurities at time of manufacture (ppb):*  
Ag 2 Cs <0.1 Ir <0.1 Pd <0.1 Tb <0.1 Al <0.1 Cu <0.1 K 15 Pt <0.1 Te <0.1 As <0.1 Dy <0.1 La <0.1 Rb <0.1 Th <0.1 Au <0.1 Er <0.1 Li <0.1 Re <0.1 Ti 4  
B <0.1 Eu <0.1 Lu <0.1 Rh <0.1 Tl <0.1 Ba <0.1 Fe <6 Mg <0.1 Ru <0.1 Tm <0.1 Be <0.1 Ga <0.1 Mn <0.1 Sb <0.1 U <0.1 Bi <0.1 Gd <0.1 Mo <0.1 Sc <0.1 V <0.1 Ca <0.1 Ge <0.1 Na 16 Se 2 W 1  
Cd <0.1 Hf <0.1 Nb 0.2 Sm <0.1 Y <0.1 Ce <0.1 Hg 3 Nd <0.1 Sn <0.1 Yb <0.1 Co <0.1 Ho <0.1 Ni <0.1 Sr <0.1 Zn 2  
Cr <0.1 In <0.1 Pb <0.1 Ta 1 Zr 2

## ROMIL-SpA™ Super Purity Acids and Reagents Specifications

Methyl Alcohol (see Methanol)

4-Methylpentan-2-one (see Methyl iso-Butyl Ketone)

### Methanol SpA

low metals

H413

500ml H413P  
1LT H413M  
2½LT H413L  
Dgr H:225-301+311+331-370  
P:210-280F-302+352-309+310-403+235



(Methyl Alcohol)

CH<sub>3</sub>OH MW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]

Assay >99.9% Water <0.05% Residue <0.0001%

Group 1 & 2 elements typically <0.5 ppb

Group 3 to 12 (transition) elements typically <0.5 ppb

Group 13, 14, 15 elements typically <0.5-50 ppb

Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis (for organic analysis and chromatography see ROMIL Code H410 Methanol SpS)

Typical values, elemental impurities at time of manufacture (ppb):

Ag	<0.5	Cd	<0.5	Mg	<0.5	Sn	<0.5
Al	<0.5	Co	<0.5	Mn	<0.5	Sr	<0.5
As	<0.5	Cr	<0.5	Mo	<0.5	Th	<0.5
B	29	Cu	<0.5	Na	<0.5	Ti	<0.5
Ba	<0.5	Fe	<0.5	Ni	<0.5	U	<0.5
Be	<0.5	Hg	<0.5	Pb	<0.5	V	<0.5
Bi	<0.5	K	<0.5	Sb	1	Zn	<0.5
Ca	<0.5	Li	<0.5	Se	<0.5	Zr	<0.5

### Methyl iso-Butyl Ketone SpA

H439

500ml H439P  
2½LT H439L  
Dgr H:225-332-319-336-351-  
EUH066  
P:210-305+351+338-304+340



(4-Methylpentan-2-one)

(CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>COCH<sub>3</sub> MW 100.16 BP 117.4°C d 0.80 CAS [108-10-1]

Assay >99.8% Water <0.01% Residue <0.0001%

Group 1 & 2 elements typically <0.1-250 ppb

Group 3 to 12 (transition) elements typically <0.1-100 ppb

Group 13, 14, 15 elements typically <0.5-100 ppb

Peroxides (at time of manufacture) <0.0001% (<1 ppm)

Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis (for organic analysis and chromatography see ROMIL Code H446 Methyl iso-Butyl Ketone SpS)

### Nitric Acid SpA

500ml H566P  
1LT H566M  
2½LT H566L  
Dgr H:272-290-331-314-EUH071  
P:260c-280c-301+330+331-305+351+338-309+310



HNO<sub>3</sub> MW 63.01 d 1.42CAS [7697-37-2] Assay 67-69%

Colour <10 Hazen (APHA)

Ag, Au, Ba, Be, Bi, Ce, Cs, Dy, Er, Eu, Ga, Gd, Ge, Hf, Hg, Ho, In, La, Li, Lu, Mn, Mo,

Nb, Nd, Pb, Pr, Rb, Re, Sc, Sm, Sr, Tb, Te, Th, Tl, Tm, U, W, Y, Yb, Zr <0.1 ppb each

As, Cd, Co, Cu, Ni, Pd, Pt, Rh, Ru, Sb, Sn, Ti, V, Zn <0.5 ppb each

Al, B, Ca, Cr, Fe, K, Mg, Na, Se <1 ppb each

Total P <0.01 ppm

Cl <0.2 ppm

Total S <0.3 ppm

Store in dark.

Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry

Elemental impurities specified at time of manufacture.

Concentrated Nitric Acid can decompose to nitrogen oxides (NO<sub>x</sub>) through action of heat or light resulting in a yellow colouration. However, this does not affect the performance of the acid with respect to trace metals or oxidising power.

Storage in a cool, dark place is recommended.

Typical values, elemental impurities at time of manufacture (ppb):

Ag <0.1 Cs <0.1 K <0.2 Pr <0.1 Tb <0.1

Al <0.5 Cu <0.1 La <0.1 Pt <0.1 Te <0.1

As <0.1 Dy <0.1 Li <0.1 Rb <0.1 Th <0.1

Au <0.1 Er <0.1 Lu <0.1 Re <0.1 Ti <0.1

B <0.5 Eu <0.1 Mg <0.2 Rh <0.1 Tl <0.1

Ba <0.1 Fe <0.5 Mn <0.1 Ru <0.1 Tm <0.1

Be <0.1 Ga <0.1 Mo <0.1 Sb <0.1 U <0.1

Bi <0.1 Gd <0.1 Na <0.2 Sc <0.1 V <0.1

Ca <0.5 Ge <0.1 Nb <0.1 Se <0.1 W <0.1

Cd <0.1 Hf <0.1 Nd <0.1 Sm <0.1 Y <0.1

Ce <0.1 Hg <0.02 Ni <0.1 Sn <0.1 Yb <0.1

Co <0.1 Ho <0.1 Pb <0.1 Sr <0.1 Zn <0.2

Cr <0.5 In <0.1 Pd <0.1 Ta <0.1 Zr <0.1

## ROMIL-SpA™ Super Purity Acids and Reagents Specifications

### Perchloric Acid SpA

less than 50% w/w

H675

500ml H675P  
2½LT H675L  
Dgr H:272-314  
P:210-220c-280c-301+330+331-  
305+351+338-309+310



HClO<sub>4</sub> MW 100.46 d 1.40 CAS [7601-90-3] □ Assay 46-49%  
Colour <10 Hazen (APHA)  
As, Au, Be, Bi, Ce, Co, Cs, Cu, Dy, Er, Eu, Ga, Gd, Ho, In, La, Li, Lu, Mo, Nd, Pd, Pr,  
Pt, Rb, Rh, Sb, Sc, Sm, Sr, Tb, Te, Tl, Tm, U, V, Y, Yb, Zr <0.5 ppb each  
Ag, Al, Ba, Ca, Cd, Fe, K, Mg, Mn, Na, Ni, Pb, Sn, Th, Ti, Zn <1 ppb each  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry

Elemental impurities specific at time of manufacture.  
Less hazardous acid concentration to facilitate international shipping.

### Perchloric Acid SpA

H584

500ml H584P  
2½LT H584L  
Dgr H:271-290-314  
P:210-221-280c-301+330+331-  
305+351+338-309+310



HClO<sub>4</sub> MW 100.46 d1.66CAS[7601-90-3] □ Assay 65-71%  
Colour <10 Hazen (APHA)  
As, Au, Be, Bi, Ce, Co, Cs, Cu, Dy, Er, Eu, Ga, Gd, Ho, In, La, Li, Lu, Mo, Nd, Pd, Pr,  
Pt, Rb, Rh, Sb, Sc, Sm, Sr, Tb, Te, Tl, Tm, U, V, Y, Yb, Zr <0.5 ppb each  
Ag, Al, Ba, Ca, Cd, Fe, K, Mg, Mn, Na, Ni, Pb, Sn, Th, Ti, Zn <1 ppb each  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry

Elemental impurities specific at time of manufacture.  
Typical values, elemental impurities at time of manufacture (ppb):  
Ag <0.1 Cu <0.1 Li <0.1 Pt <0.5 Ti <0.5 Al <0.5 Dy <0.1 Lu <0.1 Rb <0.1 Tl  
<0.1 As <0.1 Er <0.1 Mg <0.1 Rh <0.1 Tm <0.1 Au <0.1 Eu <0.1 Mn <0.1  
Sb <0.1 U <0.1 Ba <0.1 Fe <0.5 Mo <0.1 Sc <0.1 V <0.1 Be <0.1 Ga <0.1  
Na <0.1 Sm <0.1 W <0.5 Bi <0.1 Gd <0.1 Nb <0.5 Sn <0.5 Y <0.1 Ca <0.5  
Hf <0.5 Nd <0.1 Sr <0.1 Yb <0.1 Cd <0.1 Ho <0.1 Ni <0.5 Ta <0.5 Zn <0.5  
Ce <0.1 In <0.1 Pb <0.1 Tb <0.1 Zr <0.1 Co <0.1 K <0.1 Pd <0.5 Te <0.1  
Cs <0.1 La <0.1 Pr <0.1 Th <0.1

### Sulphuric Acid SpA

H691

500ml H691P  
1LT H691M  
2½LT H691L  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-  
309+310



H<sub>2</sub>SO<sub>4</sub> MW 98.07 d1.84CAS[7664-93-9] □ Assay 93-98%  
Colour <10 Hazen (APHA)  
Ba, Be, Bi, Dy, Er, Eu, Ga, Gd, Hf, Ho, La, Lu, Nd, Pb, Pr, Sc, Sm, Tb, Te, Th, Tm, U  
<0.1 ppb each  
Au, Cd, Ce, Co, Cr, Cs, Cu, In, Li, Mn, Mo, Nb, Ni, Pt, Rb, Rh, Sr, Ti, V, W, Y, Yb, Zr  
<0.5 ppb each  
Ag, Al, As, Ca, Fe, Ge, Hg, K, Mg, Na, Sb, Sn, Ti, Zn <1 ppb each  
Se <10 ppb  
Total P <0.05 ppm  
NO<sub>3</sub> <0.2 ppm  
Cl <0.7 ppm  
Substances reducing permanganate <20 ppm  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis, Ion Chromatography, Voltammetry  
Elemental impurities specific at time of manufacture.  
Typical values, elemental impurities at time of manufacture (ppb):  
Ag <0.1 Cu <0.1 La <0.1 Pt <0.5 Ti <0.5 Al <0.5 Dy <0.1 Li <0.1 Rb <0.5 Tl  
<0.5 As <1 Er <0.1 Lu <0.1 Rh <0.5 Tm <0.1 Au <0.1 Eu <0.1 Mg <0.5 Sb  
<1 U <0.1 Ba <0.1 Fe <0.5 Mn <0.5 Sc <0.1 V <0.1 Be <0.1 Ga <0.1 Mo  
<0.1 Se <5 W <0.5 Bi <0.1 Gd <0.1 Na <0.5 Sm <0.1 Y <0.5 Ca <0.5 Ge  
<0.1 Nb <0.5 Sn <0.1 Yb <0.5 Cd <0.1 Hf <0.1 Nd <0.1 Sr <0.1 Zn <0.2 Ce  
<0.5 Hg <1 Ni <0.1 Ta <0.5 Zr <0.1 Co <0.1 Ho <0.1 Pb <0.1 Tb <0.1  
Cr <0.1 In <0.5 Pd <0.5 Te <0.1  
Cs <0.5 K <0.5 Pr <0.1 Th <0.1

### Water SpA

H951

2½LT H951L

H<sub>2</sub>OMW 18.02FP 0.0°C BP 100.0°C CAS [7732-18-5] □ Residue <0.0001%  
Resistivity (at time of manufacture) >18 MOhm @ 25°C  
Elemental impurities at time of manufacture:  
Ag, Ba, Be, Bi, Cd, Hg, Mo, Pb, Sb <0.5 ppb each  
Al, As, Co, Cr, Cu, Li, Mn, Ni, Sn, Sr, Th, Ti, U, V, Zr <1 ppb each  
Fe, Mg, Zn <2 ppb each  
B <10 ppb  
Ca, K, Na <20 ppb each  
Equivalent to ASTM D1193 Type II  
Application: Environment Analysis (eg, using AAS, ICP-OES, ICP-MS), Trace Metal Analysis (for HPLC and organic trace analysis see ROMIL Code H950 Water SpS), Ion Chromatography, Voltammetry

Typical values, elemental impurities at time of manufacture (ppb):  
Ag <0.3 Ca <1 K <1 Pb <0.02  
Al <0.2 Cd <0.01 Li <0.03 Sb <0.01 V <0.05  
As <0.08 Co <0.02 Mg <0.2 Sn <0.03 Zn <0.2  
B <6.5 Cr <0.03 Mn <0.02 Sr <0.02 Zr <0.01  
Ba <0.06 Cu <0.07 Mo <0.05 Th <0.02  
Be <0.02 Fe <0.1 Na <1 Ti <0.05  
Bi <0.02 Hg <0.01 Ni <0.05 U <0.02

## ROMIL-UpA™ Ultra Purity Acids and Reagents Specifications

Acetic Acid glacial (see Acetic Acid)

Ammonium Hydroxide solution (see Ammonia solution)

### Acetic Acid UpA

SS62

500ml SS62P  
Dgr H:226-314  
P:280c-301+330+331-305+351+338-307+310



(Acetic Acid glacial)  
CH<sub>3</sub>COOHMW 60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7] Assay >99%

Trace elemental impurities: parts per trillion level  
Batch values reported on accompanying Certificate of Analysis  
Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <50	Cu <5	La <0.1	Rb <1	Th <0.1
Al <20	Dy <0.1	Li <1	Re <0.1	Ti <10
As <20	Er <0.1	Lu <0.1	Rh <20	Tl <0.1
Ba <1	Eu <0.1	Mg <10	Ru <20	Tm <0.1
Be <5	Fe <20	Mn <1	Sb <10	U <0.1
Bi <1	Ga <1	Mo <10	Sc <1	V <1
Ca <50	Gd <0.1	Na <50	Se <50	W <10
Cd <1	Ge <1	Nd <0.1	Sm <0.1	Y <1
Ce <0.1	Hf <10	Ni <10	Sn <10	Yb <0.1
Co <1	Ho <0.1	Pb <2	Sr <1	Zn <20
Cr <10	In <1	Pr <0.1	Tb <0.1	Zr <10
Cs <0.1	K <30	Pt <50	Te <1	

### Ammonia solution UpA

SS72

500ml SS72P  
Dgr H:314-335  
P:280c-301+330+331-304+340-305+351+338-309+310



(Ammonium Hydroxide solution)  
NH<sub>3</sub> MW17.03 d 0.92CAS [1336-21-6]  
Assay 20-22%

Trace elemental impurities: parts per trillion level  
Batch values reported on accompanying Certificate of Analysis  
Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <1	Cu <5	La <0.1	Pt <1	Th <0.1
Al <10	Dy <0.1	Li <1	Rb <1	Ti <5
As <10	Er <0.1	Lu <0.1	Re <1	Tl <1
Au <5	Eu <0.1	Mg <5	Rh <1	Tm <0.1
Ba <5	Fe <10	Mn <5	Ru <10	U <0.1
Be <5	Ga <1	Mo <1	Sb <10	V <2
Bi <0.1	Gd <0.1	Na <10	Sc <2	W <5
Ca <10	Ge <1	Nb <1	Se <50	Y <1
Cd <1	Hf <0.1	Nd <0.1	Sm <1	Yb <0.1
Ce <0.1	Hg <200	Ni <5	Sn <5	Zn <5
Co <1	Ho <0.1	Pb <2	Sr <1	Zr <1
Cr <5	In <1	Pd <20	Tb <0.1	
Cs <0.1	K <10	Pr <0.1	Te <1	

### Hydrobromic Acid UpA

SS82

500ml SS82P  
Dgr H:314-335  
P:280c-301+330+331-304+340-305+351+338-309+310



HBrMW80.91d 1.48CAS[10035-10-6]  
Assay 44-49%

Trace elemental impurities: parts per trillion level  
Batch values reported on accompanying Certificate of Analysis  
Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <2	Cu <5	Lu <0.01	Re <5	Tl <0.1
Al <50	Dy <0.01	Mg <5	Rh <1	Tm <0.01
Au <1	Er <0.01	Mn <2	Ru <10	U <0.01
B <1000	Eu <0.01	Mo <10	Sb <50	V <1
Ba <1	Fe <50	Na <30	Sc <1	W <10
Be <5	Ga <10	Nb <1	Sm <0.01	Y <1
Bi <0.1	Gd <0.01	Nd <0.05	Sn <20	Yb <0.01
Ca <50	Hf <0.05	Ni <10	Sr <1	Zn <5
Cd <1	Ho <0.01	Pb <1	Ta <20	Zr <1
Ce <0.05	In <1	Pd <10	Tb <0.01	
Co <1	K <20	Pr <0.05	Te <10	
Cr <10	La <0.05	Pt <1	Th <0.05	
Cs <0.05	Li <1	Rb <1	Ti <10	

### Hydrochloric Acid UpA

SS42

500ml SS42P  
Dgr H:290-314-335  
P:280c-301+330+331-305+351+338-309+310



HCIMW36.46d 1.18CAS[7647-01-0]  
Assay 32-35%

Trace elemental impurities: parts per trillion level  
Batch values reported on accompanying Certificate of Analysis  
Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <5	Cs <0.05	La <0.05	Pt <1	Te <1
Al <10	Cu <3	Li <1	Rb <1	Th <0.05
As <20	Dy <0.01	Lu <0.01	Re <0.1	Ti <10
Au <10	Er <0.01	Mg <5	Rh <1	Tl <0.1
B <10	Eu <0.01	Mn <2	Ru <10	Tm <0.01
Ba <1	Fe <10	Mo <5	Sb <20	U <0.01
Be <5	Ga <1	Na <5	Sc <1	V <1
Bi <0.05	Gd <0.01	Nb <1	Se <50	Y <0.1
Ca <10	Hf <0.05	Nd <0.05	Sm <0.01	Yb <0.01
Cd <0.1	Hg <20	Ni <10	Sn <10	Zn <5
Ce <0.05	Ho <0.01	Pb <1	Sr <1	Zr <1
Co <3	In <0.1	Pd <10	Ta <20	
Cr <10	K <10	Pr <0.05	Tb <0.01	

## ROMIL-UpA™ Ultra Purity Acids and Reagents Specifications

### Hydro uoric Acid UpA

SS52

500ml SS52P  
Dgr H:300+310+330-314  
P:280-301+330+331-302+352-  
304+340-305+351+338-  
310+ROP 010



HFMW20.01d 1.16CAS[7664-39-3]

Assay 47-51%

Trace elemental impurities: parts per trillion level

Batch values reported on accompanying Certificate of Analysis

Application: Ultra trace inorganic analysis

For treatment of HF burns, calcium gluconate gel (ROMIL Code PCG9V) is recommended. In order to provide emergency first aid, it should be kept wherever HF is handled or stored.

Typical values, elemental impurities at time of manufacture (ppt):

Ag <1 Cs <0.5 K <10 Pr <0.1 Tb <0.1  
Al <10 Cu <10 La <0.1 Pt <10 Te <1  
As <10 Dy <0.1 Li <1 Rb <1 Th <0.1  
Au <10 Er <0.1 Lu <0.1 Re <0.1 Ti <10  
B <10 Eu <0.1 Mg <5 Rh <1 Tl <0.1  
Ba <5 Fe <10 Mn <1 Ru <1 Tm <0.1  
Be <5 Ga <1 Mo <5 Sb <10 U <0.1  
Bi <0.1 Gd <0.1 Na <10 Sc <1 V <1  
Ca <10 Ge <1 Nb <5 Se <50 W <10  
Cd <0.1 Hf <1 Nd <0.1 Sm <0.1 Y <0.5  
Ce <0.1 Hg <20 Ni <10 Sn <10 Yb <0.1  
Co <1 Ho <0.1 Pb <1 Sr <1 Zn <5  
Cr <10 In <0.1 Pd <10 Ta <20 Zr <10

Typical values, anionic impurities at time of manufacture (ppb):

TotalS <50

### Hydrogen Peroxide UpA

SS92

500ml SS92P  
Dgr H:302-318  
P:280e-305+351+338-313



H2O2 MW34.01 d 1.10CAS [7722-84-1]

Assay 30-32%

Trace elemental impurities: parts per trillion level

Batch values reported on accompanying Certificate of Analysis

Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <10 Cs <1 K <10 Pr <0.5	Te <1
Al <10 Cu <5 La <0.5 Rb <5	Th <0.5
As <10 Dy <0.5 Li <1 Re <5	Ti <10
Au <10 Er <0.5 Lu <0.5 Rh <5	Tl <1
B <100 Eu <0.5 Mg <10 Ru <10	Tm <0.5
Ba <5 Fe <20 Mn <5 Sb <1	U <0.5
Be <5 Ga <5 Mo <5 Sc <10	V <5
Bi <10 Gd <0.5 Na <10 Se <50	W <10
Ca <50 Ge <10 Nb <1 Sm <0.5	Y <1
Cd <1 Hf <1 Nd <0.5 Sn <10	Yb <0.5
Ce <1 Hg <20 Ni <10 Sr <1	Zn <10
Co <5 Ho <0.5 Pb <1 Ta <10	Zr <5
Cr <5 In <0.5 Pd <10 Tb <0.5	

### Nitric Acid UpA

SS12

500ml SS12P  
Dgr H:272-290-331-314-EUH071  
P:260c-280c-301+330+331-  
305+351+338-309+310



HNO3 MW63.01 d1.42CAS [7697-37-2]

Assay 67-69%

Trace elemental impurities: parts per trillion level

Batch values reported on accompanying Certificate of Analysis

Store in dark.

Application: Ultra trace inorganic analysis

Concentrated Nitric Acid can decompose nitrogen oxides (NOx) through action of heat or light resulting in a yellow colouration. However, this does not affect the performance of the acid with respect to trace metals or oxidising power.

Storage in a cool, dark place is recommended.

Typical values, elemental impurities at time of manufacture (ppt):

Ag <2 Cs <0.05 K <5 Pr <0.05 Tb <0.01  
Al <10 Cu <3 La <0.05 Pt <1 Te <1  
As <10 Dy <0.01 Li <1 Rb <1 Th <0.05  
Au <10 Er <0.01 Lu <0.01 Re <1 Ti <10  
B <10 Eu <0.01 Mg <5 Rh <1 Tl <0.1  
Ba <1 Fe <10 Mn <2 Ru <10 Tm <0.01  
Be <5 Ga <1 Mo <1 Sb <10 U <0.01  
Bi <0.1 Gd <0.01 Na <5 Sc <1 V <1  
Ca <10 Ge <1 Nb <1 Se <20 W <5  
Cd <1 Hf <0.05 Nd <0.05 Sm <0.01 Y <1  
Ce <0.05 Hg <20 Ni <10 Sn <10 Yb <0.01  
Co <1 Ho <0.01 Pb <1 Sr <1 Zn <5  
Cr <10 In <1 Pd <10 Ta <10 Zr <1

### Perchloric Acid UpA

SS22

500ml SS22P  
Dgr H:271-290-314  
P:210-221-280c-301+330+331-  
305+351+338-309+310



HClO4 MW100.46 d1.66CAS[7601-90-3]

Assay 65-71%

Trace elemental impurities: parts per trillion level

Batch values reported on accompanying Certificate of Analysis

Application: Ultra trace inorganic analysis

## ROMIL-UpA™ Ultra Purity Acids and Reagents Specifications

### Sulphuric Acid UpA

SS32

 500ml SS32P  
 Dgr H:290-314  
 P:280c-301+330+331-305+351+338-  
 309+310


H<sub>2</sub>SO<sub>4</sub> MW98.07 d1.84CAS [7664-93-9] □

Assay 93-98%

Trace elemental impurities: parts per trillion level

Batch values reported on accompanying Certi cate of Analysis

Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <5	Dy <0.1	Li <10	Rb <1	Tl <1
Al <30	Er <0.1	Lu <0.1	Rh <1	Tm <0.1
As <500	Eu <0.1	Mg <10	Sb <10	U <0.1
Ba <10	Fe <30	Mn <1	Sc <5	V <5
Be <5	Ga <1	Mo <10	Se <500	W <5
Bi <1	Gd <0.1	Na <30	Sm <0.1	Y <1
Ca <50	Ge <100	Nb <1	Sn <50	Yb <0.1
Cd <1	Hf <0.1	Nd <0.1	Sr <1	Zn <20
Ce <0.1	Hg <50	Ni <20	Ta <20	Zr <5
Co <1	Ho <0.1	Pb <5	Tb <0.1	
Cr <10	In <1	Pd <10	Te <0.1	
Cs <1	K <50	Pr <0.1	Th <0.1	
Cu <5	La <1	Pt <10	Ti <50	

### Water UpA

SS02

1LT SS02M

H<sub>2</sub>O MW18.02 FP0.0°C BP 100.0°C CAS [7732-18-5] □

Trace elemental impurities: parts per trillion level

Trace anionic impurities: parts per billion level

Batch values reported on accompanying Certi cate of Analysis

Equivalent to ASTM D1193 Type I

Application: Ultra trace inorganic analysis

Typical values, elemental impurities at time of manufacture (ppt):

Ag <5	Cs <0.1	K <10	Pr <0.1	Tb <0.1
Al <20	Cu <5	La <0.1	Pt <1	Te <1
As <10	Dy <0.1	Li <1	Rb <1	Th <0.1
Au <10	Er <0.1	Lu <0.1	Re <1	Ti <10
B <20	Eu <0.1	Mg <5	Rh <1	Tl <0.1
Ba <1	Fe <10	Mn <1	Ru <1	Tm <0.1
Be <5	Ga <1	Mo <1	Sb <10	U <0.1
Bi <0.1	Gd <0.1	Na <10	Sc <1	V <1
Ca <10	Ge <1	Nb <1	Se <50	W <10
Cd <1	Hf <0.1	Nd <0.1	Sm <0.1	Y <1
Ce <0.1	Hg <20	Ni <10	Sn <10	Yb <0.1
Co <1	Ho <0.1	Pb <1	Sr <1	Zn <10
Cr <10	In <1	Pd <5	Ta <10	Zr <1

Typical values, anionic impurities at time of manufacture (ppb):

Cl <1 PO<sub>4</sub> <1 SO<sub>4</sub> <1



## ROMIL High Purity Aqueous Mixes

for instrumental techniques

## ROMIL-SpA™ Ion Chromatography Solutions Specifications

IC Eluant, Regenerant & Concentrate mixes

### IC Concentrate CHC1 SpA

carbonate/hydrogen carbonate

RJ183

1LT RJ183M  
2½LT RJ183L

3 components:  
Water (ROMIL H951)  
containing  
Sodium Carbonate anhydrous 64 mM  
Sodium Hydrogen Carbonate 20 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Eluant concentrate (20x) for ion chromatography, Metrosep A Supp 5

### IC Concentrate DPAN1 SpA

dipicolinic acid/nitric acid

RJ239

1LT RJ239M  
2½LT RJ239L

3 components:  
Water (ROMIL H951)  
containing  
Dipicolinic Acid 17 mM  
Nitric Acid 17 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Eluant concentrate (10x) for ion chromatography, Metrosep C 6

### IC Concentrate DPAN2 SpA

dipicolinic acid/nitric acid

RJ328

1LT RJ328M  
2½LT RJ328L

3 components:  
Water (ROMIL H951)  
containing  
Dipicolinic Acid 14 mM  
Nitric Acid 34 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Eluant concentrate (20x) for ion chromatography, Metrosep C 4

### IC Concentrate mix PHSF1 SpA

pdca/hydroxide/sulphate/formic acid

RJ812

1LT RJ812M  
2½LT RJ812L  
Wng H:315-319  
P:302+352-305+351+338

5 components:  
Water (ROMIL H951)  
containing  
Pyridine-2,6-dicarboxylic acid (PDCA) 35 mM  
Potassium Hydroxide 330 mM  
Potassium Sulphate 28 mM  
Formic Acid 370 mM  
pH after dilution 4.2 ±0.1  
Protect from atmospheric CO<sub>2</sub>.  
Application: Eluant concentrate (5x) for ion chromatography, equivalent to Dionex 046088  
*Dionex is a registered trademark of Dionex Corporation*



### IC Diluent mix DHHC1 SpA

dmae/hydroxide/hydrogen carbonate

RJ407

1LT RJ407M  
2½LT RJ407L  
Dgr H:314-318-335-412  
P:280c-260-273-303+361+353-305+351+338-304+340-312

4 components:  
Water (ROMIL H951)  
containing  
2-Dimethylaminoethanol (DMAE) 1 M  
Ammonium Hydroxide 0.5 M  
Sodium Hydrogen Carbonate 0.3 M  
pH 10.4 ±0.2  
Absorbance after addition of 4-(2-pyridylazo)resorcinol (PAR) 0.5 mM:  
530nm <0.5 AU  
Protect from atmospheric CO<sub>2</sub>.  
Application: Postcolumn diluent solution for ion chromatography for detection of transition metals, equivalent to Dionex 046094  
*Dionex is a registered trademark of Dionex Corporation*



### IC Eluant C2 SpA

sodium carbonate 3.6 mM

RJ924

1LT RJ924M  
2½LT RJ924L

2 components:  
Water (ROMIL H951)  
containing  
Sodium Carbonate 3.6 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Eluant solution for ion chromatography

## ROMIL High Purity Aqueous Mixes

for instrumental techniques

### ROMIL-SpA™ Ion Chromatography Solutions Specifications

IC Eluant, Regenerant & Concentrate mixes

#### IC Eluant N2 SpA

nitric acid 7.25 mM

RJ701

1LT RJ701M  
2½LT RJ701L

2 components:  
Water (ROMIL H951)  
containing  
Nitric Acid 7.25 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Eluant solution for ion chromatography, Metrosep Fast C6

#### IC Regenerant CHC2 SpA

carbonate/hydrogen carbonate

RJ511

1LT RJ511M  
2½LT RJ511L

3 components:  
Water (ROMIL H951)  
containing  
Sodium Carbonate anhydrous 70 mM  
Sodium Hydrogen Carbonate 70 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Cation suppressor regenerant solution for ion chromatography

#### IC Regenerant OS1 SpA

oxalate/sulphuric acid

RJ610

1LT RJ610M  
2½LT RJ610L

3 components:  
Water (ROMIL H951)  
containing  
Oxalic Acid 100 mM  
Sulphuric Acid 500 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Anion suppressor regenerant solution for ion chromatography, extends resin life by minimising chelation of transition metals

#### IC Regenerant RN1 SpA

rubidium/nitric acid

RJ306

1LT RJ306M  
2½LT RJ306L

3 components:  
Water (ROMIL H951)  
containing  
Rubidium Nitrate as Rb 1 mg/LT  
Nitric Acid 100 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Cation suppressor regenerant concentrate (20x) for ion chromatography, Metrosep C Supp 1 & 2

#### IC Wash solution E2 SpA

edta na2 3 mM

RJ623

500ml RJ623P  
1LT RJ623M

2 components:  
Water (ROMIL H951)  
containing  
EDTA di-Sodium salt 3 mM  
Protect from atmospheric CO<sub>2</sub>.  
Application: Wash solution for ion chromatography

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Acetic Acid 1M PrimAg-TvR

V1010

1LT V1010M CH<sub>3</sub>COOH MW60.05  
 2½LT V1010L Nominal concentration 1M (1N) ±0.5% @ 20°C  
 5LT V1010K CH<sub>3</sub>COOH 60.05 g/LT  
 10LT V1010J Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*

### Acetic Acid 0.05M/0.1M/0.2M PrimAg-TvR tri-concentrate

C2062

6x100ml C2062Q CH<sub>3</sub>COOH MW 60.05  
 1M (1N) nominal concentrate ±0.5% @ 20°C  
 CH<sub>3</sub>COOH 60.05 g/LT  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 2.0LT yields 0.05M (0.05N)  
 Dilute to 1.0LT yields 0.1M (0.1N)  
 Dilute to 0.5LT yields 0.2M (0.2N)  
 Application: Concentrate for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*  
 Pack contains 6 vials each containing 100ml of concentrate.

### Ammonia 5M PrimAg-TvR in ethanol/water 69:31

V1205

2½LT V1205L NH<sub>3</sub> MW 17.03  
 Dgr H:225-314-335 Nominal concentration 5M (5N) ±0.5% @ 20°C  
 P:210-280c-304+340-305+351+338- NH<sub>3</sub> 85.15 g/LT  
 309+310 Solvent: CH<sub>3</sub>CH<sub>2</sub>OH/H<sub>2</sub>O 69:31 v/v  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Ammonium Cerium(IV) Sulphate 0.1M PrimAg-TvR

V0218

1LT V0218M (NH<sub>4</sub>)<sub>4</sub>Ce(SO<sub>4</sub>)<sub>4</sub> MW596.52  
 2½LT V0218L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 Wng H:290-315-319 (NH<sub>4</sub>)<sub>4</sub>Ce(SO<sub>4</sub>)<sub>4</sub> 59.65 g/LT  
 P:302+352-305+351+338 H<sub>2</sub>SO<sub>4</sub> 150 g/LT  
 Application: Stable alternative to potassium permanganate in oxidation titrimetry  
*ISO17034 accredited SI-traceable*



### Ammonium Hydroxide 5M PrimAg-TvR

V3297

1LT V3297M NH<sub>4</sub>OH MW35.05  
 2½LT V3297L Nominal concentration 5M (5N) ±0.5% @ 20°C  
 Dgr H:314-335 NH<sub>4</sub>OH 175.3g/LT  
 P:280c-301+330+331-304+340- Application: Reagent for volumetric titrimetry  
 305+351+338-309+310 *ISO17034 accredited SI-traceable*



### di-Ammonium Iron(II) Sulphate 0.025M PrimAg-TvR

V0224

1LT V0224M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub> MW284.01  
 2½LT V0224L Nominal concentration 0.025M (0.025N)  
 (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub> 7.100 g/LT  
 Stabiliser: H<sub>2</sub>SO<sub>4</sub> 5 g/LT  
 Calibrate by titration with either Ce(SO<sub>4</sub>)<sub>2</sub>, K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> or KMnO<sub>4</sub> of equivalent molarity immediately before each use.  
 Application: Reagent for volumetric titrimetry

### di-Ammonium Iron(II) Sulphate 0.1M PrimAg-TvR

V0242

1LT V0242M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub> MW284.01  
 2½LT V0242L Nominal concentration 0.1M (0.1N)  
 (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub> 28.40 g/LT  
 Stabiliser: H<sub>2</sub>SO<sub>4</sub> 5 g/LT  
 Calibrate by titration with either Ce(SO<sub>4</sub>)<sub>2</sub>, K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> or KMnO<sub>4</sub> of equivalent molarity immediately before each use.  
 Application: Reagent for volumetric titrimetry

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### di-Ammonium Iron(II) Sulphate 0.3M

#### PrimAg-TvR

V0235

1LT V0235M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub> MW284.01  
 2½LT V0235L Nominal concentration 0.3M (0.3N)  
 (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub> 85.20 g/LT  
 Stabiliser: H<sub>2</sub>SO<sub>4</sub> 5 g/LT  
 Calibrate by titration with either Ce(SO<sub>4</sub>)<sub>2</sub>, K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> or KMnO<sub>4</sub> of equivalent molarity immediately before each use.  
 Application: Reagent for volumetric titrimetry

### Ammonium Thiocyanate 0.1M PrimAg-TvR

V2703

1LT V2703M NH<sub>4</sub>SCN MW76.12  
 2½LT V2703L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 5LT V2703K NH<sub>4</sub>SCN 7.612 g/LT  
 10LT V2703J Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

**Benzethonium Chloride \*M (see Hyamine 1622 \*M)**

### Barium Hydroxide 0.05M PrimAg-TvR

V1279

1LT V1279M Ba(OH)<sub>2</sub> MW171.34  
 2½LT V1279L Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C  
 Ba(OH)<sub>2</sub> 8.567 g/LT  
 Manufactured and filled under N<sub>2</sub>.  
 Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Calcium Chloride 0.25M PrimAg-TvR

V3191

1LT V3191M CaCl<sub>2</sub> MW110.99  
 2½LT V3191L Nominal concentration 0.25M (0.5N) ±0.5% @ 20°C  
 5LT V3191K CaCl<sub>2</sub> 27.75 g/LT  
 10LT V3191J Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Calcium Chloride 1M PrimAg-TvR

V2182


1LT V2182M CaCl<sub>2</sub> MW110.99  
 2½LT V2182L Nominal concentration 1M (2N) ±0.5% @ 20°C  
 5LT V2182K CaCl<sub>2</sub> 111.0 g/LT  
 10LT V2182J Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Cerium(IV) Sulphate 0.05M/0.1M

#### PrimAg-TvR

di-concentrate

C3875

6x100ml C3875Q Ce(SO<sub>4</sub>)<sub>2</sub> MW 332.22  
 Dgr H:290-314 0.5M (0.5N) nominal concentrate ±0.5% @ 20°C  
 P:280c-301+330+331-305+351+338- Ce(SO<sub>4</sub>)<sub>2</sub> 166.1 g/LT  
 309+310 H<sub>2</sub>SO<sub>4</sub> 200 g/LT  
  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 1.0LT yields 0.05M (0.05N)  
 Dilute to 0.5LT yields 0.1M (0.1N)  
 Application: Stable alternative to potassium permanganate in oxidation titrimetry


ISO 17034 accredited SI-traceable

Pack contains 6 vials each containing 100ml of concentrate.

### Cerium(IV) Sulphate 0.1M PrimAg-TvR

in sulphuric acid 4%


V2764

1LT V2764M Ce(SO<sub>4</sub>)<sub>2</sub> MW 332.22  
 2½LT V2764L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 Wng H:290-315-319 Ce(SO<sub>4</sub>)<sub>2</sub> 33.22 g/LT  
 P:302+352-305+351+338 H<sub>2</sub>SO<sub>4</sub> 40 g/LT  
  
 Application: Stable alternative to potassium permanganate in oxidation titrimetry  
 ISO17034 accredited SI-traceable

### Cerium(IV) Sulphate 0.1M PrimAg-TvR

in sulphuric acid 10%

V4582

1LT V4582M Ce(SO<sub>4</sub>)<sub>2</sub> MW332.22  
 2½LT V4582L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 Wng H:290-315-319 Ce(SO<sub>4</sub>)<sub>2</sub> 33.22 g/LT  
 P:302+352-305+351+338 H<sub>2</sub>SO<sub>4</sub> 100 g/LT  
  
 Application: Stable alternative to potassium permanganate in oxidation titrimetry  
 ISO17034 accredited SI-traceable

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Dodecylbenzenesulphonic Acid 0.002M

#### PrimAg-TvR

sodium salt

V2543

1LT V2543M	(Sodium Dodecylbenzenesulphonate 0.002M)
2½LT V2543L	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>11</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>3</sub> .Na MW348.48
	Nominal concentration 0.002M ±0.5% @ 20°C
	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>11</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>3</sub> .Na 0.6970 g/LT
	Application: Determination of cationic surfactants
	ISO17034 accredited SI-traceable

### EDTA di-Sodium salt 0.01M PrimAg-TvR

V2934

1LT V2934M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V2934L	Nominal concentration 0.01M (0.02N) ±0.5% @ 20°C
5LT V2934K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 3.362 g/LT
10LT V2934J	Application: Reagent for volumetric titrimetry
	ISO 17034 accredited SI-traceable

### EDTA di-Sodium salt 0.02M PrimAg-TvR

V1823

1LT V1823M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V1823L	Nominal concentration 0.02M (0.04N) ±0.5% @ 20°C
5LT V1823K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 6.724 g/LT
10LT V1823J	Application: Reagent for volumetric titrimetry
	ISO 17034 accredited SI-traceable

### EDTA di-Sodium salt 0.01M/0.02M/0.04M

#### PrimAg-TvR

tri-concentrate

C5263

6x100ml C5263Q	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
	0.2M (0.4N) nominal concentration ±0.5% @ 20°C
	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 67.24 g/LT
	Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:
	Dilute to 2.0LT yields 0.01M (0.02N)
	Dilute to 1.0LT yields 0.02M (0.04N)
	Dilute to 0.5LT yields 0.04M (0.08N)
	Application: Concentrate for volumetric titrimetry
	ISO 17034 accredited SI-traceable
	Pack contains 6 vials each containing 100ml of concentrate.

### EDTA di-Sodium salt 0.05M PrimAg-TvR

V3623

1LT V3623M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V3623L	Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C
5LT V3623K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 16.81 g/LT
10LT V3623J	Application: Reagent for volumetric titrimetry
	ISO 17034 accredited SI-traceable

### EDTA di-Sodium salt 0.1M PrimAg-TvR

V3045

1LT V3045M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V3045L	Nominal concentration 0.1M (0.2N) ±0.5% @ 20°C
5LT V3045K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 33.62 g/LT
10LT V3045J	Application: Reagent for volumetric titrimetry
	ISO 17034 accredited SI-traceable

### Hyamine 1622 0.004M PrimAg-TvR

V7846

1LT V7846M	(Benzethonium Chloride 0.004M)
2½LT V7846L	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl MW448.08
Wng H:315-319-411	Nominal concentration 0.004M
P:273-302+352-305+351+338	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl 1.792 g/LT
	Calibrate by titration with Sodium Dodecyl Sulphate immediately before each use.
	Application: Determination of anionic surfactants
	Hyamine is a registered trademark of
	Lonza Inc



### Hyamine 1622 0.04M PrimAg-TvR

V6236

1LT V6236M	(Benzethonium Chloride 0.04M)
2½LT V6236L	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl MW448.08
Wng H:315-319-411	Nominal concentration 0.04M
P:273-302+352-305+351+338	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl 17.92 g/LT
	Calibrate by titration with Sodium Dodecyl Sulphate immediately before each use.
	Application: Determination of anionic surfactants
	Hyamine is a registered trademark of Lonza Inc



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Dodecylbenzenesulphonic Acid 0.002M V2543

#### PrimAg-TvR

##### sodium salt

1LT V2543M	(Sodium Dodecylbenzenesulphonate 0.002M)
2½LT V2543L	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>11</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>3</sub> .Na MW348.48
	Nominal concentration 0.002M ±0.5% @ 20°C
	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>11</sub> C <sub>6</sub> H <sub>4</sub> SO <sub>3</sub> .Na 0.6970 g/LT
	Application: Determination of cationic surfactants
	<i>ISO17034 accredited SI-traceable</i>

### EDTA di-Sodium salt 0.01M PrimAg-TvR V2934

1LT V2934M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V2934L	Nominal concentration 0.01M (0.02N) ±0.5% @ 20°C
5LT V2934K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 3.362 g/LT
10LT V2934J	Application: Reagent for volumetric titrimetry
	<i>ISO 17034 accredited SI-traceable</i>

### EDTA di-Sodium salt 0.02M PrimAg-TvR V1823

1LT V1823M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V1823L	Nominal concentration 0.02M (0.04N) ±0.5% @ 20°C
5LT V1823K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 6.724 g/LT
10LT V1823J	Application: Reagent for volumetric titrimetry
	<i>ISO 17034 accredited SI-traceable</i>

### EDTA di-Sodium salt 0.01M/0.02M/0.04M

#### PrimAg-TvR

##### tri-concentrate C5263

6x100ml C5263Q	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
	0.2M (0.4N) nominal concentration ±0.5% @ 20°C
	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 67.24 g/LT
	Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:
	Dilute to 2.0LT yields 0.01M (0.02N)
	Dilute to 1.0LT yields 0.02M (0.04N)
	Dilute to 0.5LT yields 0.04M (0.08N)
	Application: Concentrate for volumetric titrimetry
	<i>ISO 17034 accredited SI-traceable</i>
	<i>Pack contains 6 vials each containing 100ml of concentrate.</i>


### EDTA di-Sodium salt 0.05M PrimAg-TvR V3623

1LT V3623M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V3623L	Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C
5LT V3623K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 16.81 g/LT
10LT V3623J	Application: Reagent for volumetric titrimetry
	<i>ISO 17034 accredited SI-traceable</i>


### EDTA di-Sodium salt 0.1M PrimAg-TvR V3045

1LT V3045M	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> MW 336.20
2½LT V3045L	Nominal concentration 0.1M (0.2N) ±0.5% @ 20°C
5LT V3045K	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>8</sub> Na <sub>2</sub> 33.62 g/LT
10LT V3045J	Application: Reagent for volumetric titrimetry
	<i>ISO 17034 accredited SI-traceable</i>

### Hyamine 1622 0.004M PrimAg-TvR V7846

1LT V7846M	(Benzethonium Chloride 0.004M)
2½LT V7846L	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl MW448.08
Wng H315-319-411	Nominal concentration 0.004M
P:273-302+352-305+351+338	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl 1.792 g/LT
	Calibrate by titration with Sodium Dodecyl Sulphate immediately before each use.
	Application: Determination of anionic surfactants
	<i>Hyamine is a registered trademark of Lonza Inc</i>

### Hyamine 1622 0.04M PrimAg-TvR V6236

1LT V6236M	(Benzethonium Chloride 0.04M)
2½LT V6236L	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl MW448.08
Wng H315-319-411	Nominal concentration 0.04M
P:273-302+352-305+351+338	C <sub>27</sub> H <sub>42</sub> N <sub>2</sub> O <sub>2</sub> Cl 17.92 g/LT
	Calibrate by titration with Sodium Dodecyl Sulphate immediately before each use.
	Application: Determination of anionic surfactants
	<i>Hyamine is a registered trademark of Lonza Inc</i>

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Hyamine 1622 0.05M PrimAg-TvR

V4868

1LT V4868M (Benzethonium Chloride 0.05M)  
 2½LT V4868L C27H42NO2Cl MW448.08  
 Wng H:315-319-411 Nominal concentration 0.05M  
 P:273-302+352-305+351+338 C27H42NO2Cl 22.40 g/LT



Calibrate by titration with Sodium Dodecyl Sulphate immediately before each use.

Application: Determination of anionic surfactants

Hyamine is a registered trademark of Lonza Inc

### Hyamine 1622 0.1M PrimAg-TvR

V5979

1LT V5979M (Benzethonium Chloride 0.1M)  
 2½LT V5979L C27H42NO2Cl MW448.08  
 Wng H:315-319-411 Nominal concentration 0.1M  
 P:273-302+352-305+351+338 C27H42NO2Cl 44.81 g/LT



Calibrate by titration with Sodium Dodecyl Sulphate immediately before each use.

Application: Determination of anionic surfactants

Hyamine is a registered trademark of Lonza Inc

### Hydrochloric Acid 0.01M PrimAg-TvR

V3604

1LT V3604M HCl MW 36.46  
 2½LT V3604L Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 5LT V3604K HCl 0.3646 g/LT  
 10LT V3604J Application: Reagent for volumetric titrimetry

### Hydrochloric Acid 0.02M PrimAg-TvR

V3718

1LT V3718M HCl MW 36.46  
 2½LT V3718L Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 5LT V3718K HCl 0.7292 g/LT  
 10LT V3718J Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Hydrochloric Acid 0.04M PrimAg-TvR

high accuracy

V9504

1LT V9504M HCl MW 36.46  
 2½LT V9504L Nominal concentration 0.04M (0.04N) ±0.01% @ 20°C  
 Wng H:290 HCl 1.4584 g/LT  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Hydrochloric Acid 0.1M PrimAg-TvR

V3721

1LT V3721M HCl MW 36.46  
 2½LT V3721L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 5LT V3721K HCl 3.646 g/LT  
 10LT V3721J Application: Reagent for volumetric titrimetry  
 Wng H:290 ISO17034 accredited SI-traceable



### Hydrochloric Acid 0.05M/0.1M/0.2M

PrimAg-TvR

tri-concentrate

C4832

6x100ml C4832Q HCl MW 36.46  
 Wng H:290 1M (1N) nominal concentrate ±0.5% @ 20°C  
 HCl 36.46 g/LT



Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:

Dilute to 2.0LT yields 0.05M (0.05N)

Dilute to 1.0LT yields 0.1M (0.1N)

Dilute to 0.5LT yields 0.2M (0.2N)

Application: Concentrate for volumetric titrimetry

ISO 17034 accredited SI-traceable

Pack contains 6 vials each containing 100ml of concentrate.

### Hydrochloric Acid 0.2M PrimAg-TvR

V3723

1LT V3723M HCl MW 36.46  
 2½LT V3723L Nominal concentration 0.2M (0.2N) ±0.5% @ 20°C  
 5LT V3723K HCl 7.292 g/LT  
 10LT V3723J Application: Reagent for volumetric titrimetry  
 Wng H:290 ISO17034 accredited SI-traceable





## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Hydrochloric Acid 0.2M PrimAg-TvR high accuracy

V9521

1LT V9521M HCl MW 36.46  
 2½LT V9521L Nominal concentration 0.2M (0.2N) ±0.01% @ 20°C  
 Wng H:290 HCl 7.2920 g/LT  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable



### Hydrochloric Acid 0.5M PrimAg-TvR

V3725

1LT V3725M HCl MW 36.46  
 2½LT V3725L Nominal concentration 0.5M (0.5N) ±0.5% @ 20°C  
 5LT V3725K HCl 18.23 g/LT  
 10LT V3725J Application: Reagent for volumetric titrimetry  
 Wng H:290 ISO17034 accredited SI-traceable



### Hydrochloric Acid 0.25M/0.5M/1M PrimAg-TvR

tri-concentrate

C3462

6x100ml C3462Q HCl MW 36.46  
 Wng H:290-315-319-335 5M (5N) nominal concentration ±0.5% @ 20°C  
 P:302+352- HCl 182.3 g/LT  
 305+351+338 Dilution of the vial contents with distilled water using class A glassware gives  
 the following working strengths:  
 Dilute to 2.0LT yields 0.25M (0.25N)  
 Dilute to 1.0LT yields 0.5M (0.5N)  
 Dilute to 0.5LT yields 1M (1N)  
 Application: Concentrate for volumetric titrimetry  
 ISO 17034 accredited SI-traceable  
 Pack contains 6 vials each containing 100ml of concentrate.



### Hydrochloric Acid 1M PrimAg-TvR

V3772

1LT V3772M HCl MW 36.46  
 2½LT V3772L Nominal concentration 1M (1N) ±0.5% @ 20°C  
 5LT V3772K HCl 36.46 g/LT  
 10LT V3772J Application: Reagent for volumetric titrimetry  
 Wng H:290 ISO17034 accredited SI-traceable



### Hydrochloric Acid 1M PrimAg-TvR high accuracy

V9532

1LT V9532M HCl MW 36.46  
 2½LT V9532L Nominal concentration 1M (1N) ±0.01% @ 20°C  
 Wng H:290 HCl 36.460 g/LT  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable



### Hydrochloric Acid 2M PrimAg-TvR

V3775

1LT V3775M HCl MW 36.46  
 2½LT V3775L Nominal concentration 2M (2N) ±0.5% @ 20°C  
 5LT V3775K HCl 72.92 g/LT  
 10LT V3775J Application: Reagent for volumetric titrimetry  
 Wng H:290 ISO17034 accredited SI-traceable



### Hydrochloric Acid 3M PrimAg-TvR

V4758

1LT V4758M HCl MW 36.46  
 2½LT V4758L Nominal concentration 3M (3N) ±0.5% @ 20°C  
 5LT V4758K HCl 109.4 g/LT  
 10LT V4758J Application: Reagent for volumetric titrimetry  
 Wng H:290-315-319-335 ISO17034 accredited SI-traceable  
 P:302+352-305+351+338



### Hydrochloric Acid 4M PrimAg-TvR

V3736

1LT V3736M HCl MW 36.46  
 2½LT V3736L Nominal concentration 4M (4N) ±0.5% @ 20°C  
 5LT V3736K HCl 145.8 g/LT  
 10LT V3736J Application: Reagent for volumetric titrimetry  
 Wng H:290-315-319-335 ISO17034 accredited SI-traceable  
 P:302+352-305+351+338



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Hydrochloric Acid 5M PrimAg-TvR

V3777

1LT V3777M HCl MW 36.46  
 2½LT V3777L Nominal concentration 5M (5N) ±0.5% @ 20°C  
 5LT V3777K HCl 182.3 g/LT  
 10LT V3777J Application: Reagent for volumetric titrimetry  
 Wng H:290-315-319-335 *ISO17034 accredited SI-traceable*  
 P:302+352-  
 305+351+338



### Hydro uoric Acid 3M PrimAg-TvR

V5253

1LT V5253M HF MW 20.01  
 Dgr H:301+331-310-314 Nominal concentration 3M (3N) ±0.5% @ 20°C  
 P:280-301+330+331-302+352- HF 60.02 g/LT  
 304+340-305+351+338-309+310 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Iodine 0.0236M PrimAg-TvR

V3292

1LT V3292M I2 MW253.81  
 2½LT V3292L Nominal concentration 0.0236M (0.0473N) ±0.5% @ 20°C  
 Dgr H:372-401 I2 5.990 g/LT  
 P:260v-264-273-314 KI 57 g/LT  
 Store in dark.  
 Application: ASTM Iodine absorption number test  
*ISO17034 accredited SI-traceable*



### Iodine 0.025M PrimAg-TvR

V5337

1LT V5337M I2 MW253.81  
 2½LT V5337L Nominal concentration 0.025M (0.05N) ±0.5% @ 20°C  
 Dgr H:372-401 I2 6.345 g/LT  
 P:260v-264-273-314 KI 10 g/LT  
 Store in dark.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Iodine 0.05M PrimAg-TvR

V5056

1LT V5056M I2 MW253.81  
 2½LT V5056L Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C  
 Dgr H:372-401 I2 12.69 g/LT  
 P:260v-264-273-314 KI 20 g/LT  
 Store in dark.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Iodine 0.025M/0.05M/0.1M PrimAg-TvR tri-concentrate

C6167

6x100ml C6167Q I2 MW 253.81  
 Dgr H:372-401 0.5M (1N) nominal concentrate ±0.5% @ 20°C  
 P:260v-264-273-314 I2 126.9 g/LT  
 KI 200 g/LT  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 2.0LT yields 0.025M (0.05N)  
 Dilute to 1.0LT yields 0.05M (0.1N)  
 Dilute to 0.5LT yields 0.1M (0.2N)  
 Store in dark.  
 Application: Concentrate for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*  
 Pack contains 6 vials each containing 100ml of concentrate.



### Iodine 0.5M PrimAg-TvR

V4043

500ml V4043P I2 MW253.81  
 1LT V4043M Nominal concentration 0.5M (1N) ±0.5% @ 20°C  
 2½LT V4043L I2 126.9 g/LT  
 Dgr H:372-401 KI 200 g/LT  
 P:260v-264-273-314 Store in dark.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Nitric Acid 0.02M PrimAg-TvR

V5311

1LT V5311M HNO3 MW 63.01  
 2½LT V5311L Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 5LT V5311K HNO3 1.260 g/LT  
 10LT V5311J Stabiliser: Hg(NO3)2.½H2O 10 ppm  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Nitric Acid 0.02M PrimAg-TvR

V5311

1LT V5311M HNO<sub>3</sub> MW 63.01  
 2½LT V5311L Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 5LT V5311K HNO<sub>3</sub> 1.260 g/LT  
 10LT V5311J Stabiliser: Hg(NO<sub>3</sub>)<sub>2</sub>·½H<sub>2</sub>O 10 ppm  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Nitric Acid 0.1M PrimAg-TvR

V4813

1LT V4813M HNO<sub>3</sub> MW63.01  
 2½LT V4813L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 5LT V4813K HNO<sub>3</sub> 6.301 g/LT  
 10LT V4813J Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Nitric Acid 0.5M PrimAg-TvR

V3642

1LT V3642M HNO<sub>3</sub> MW63.01  
 2½LT V3642L Nominal concentration 0.5M (0.5N) ±0.5% @ 20°C  
 5LT V3642K HNO<sub>3</sub> 31.51 g/LT  
 10LT V3642J Application: Reagent for volumetric titrimetry  
 Dgr H:315-318  
*ISO17034 accreditedSI-traceable*  
 P:280e-302+352-305+351+338-310



### Nitric Acid 1M PrimAg-TvR

V4257

1LT V4257M HNO<sub>3</sub> MW63.01  
 2½LT V4257L Nominal concentration 1M (1N) ±0.5% @ 20°C  
 5LT V4257K HNO<sub>3</sub> 63.01 g/LT  
 10LT V4257J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314-EUH071  
*ISO17034 accreditedSI-traceable*  
 P:280c-302+352-305+351+338-310



### Nitric Acid 4M PrimAg-TvR

V4468

1LT V4468M HNO<sub>3</sub> MW63.01  
 2½LT V4468L Nominal concentration 4M (4N) ±0.5% @ 20°C  
 5LT V4468K HNO<sub>3</sub> 252.0 g/LT  
 10LT V4468J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314-EUH071  
*ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Nitric Acid 8M PrimAg-TvR

V5419

1LT V5419M HNO<sub>3</sub> MW63.01  
 2½LT V5419L Nominal concentration 8M (8N) ±0.5% @ 20°C  
 5LT V5419K HNO<sub>3</sub> 504.0 g/LT  
 10LT V5419J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314-EUH071  
*ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Oxalic Acid 0.05M PrimAg-TvR

V0361

1LT V0361M (COOH)<sub>2</sub> MW90.03  
 2½LT V0361L Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C  
 5LT V0361K (COOH)<sub>2</sub> 4.502 g/LT  
 10LT V0361J Protect from air.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Perchloric Acid 0.1M PrimAg-TvR in acetic acid

V6373

500ml V6373P HClO<sub>4</sub> MW 100.46  
 1LT V6373M Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 2½LT V6373L HClO<sub>4</sub> 10.05 g/LT  
 Dgr H:226-314  
 Solvent: CH<sub>3</sub>COOH  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-307+310



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Potassium Bromate/Bromide 0.1N

#### PrimAg-TvR

V7391

1LT V7391M KBrO<sub>3</sub> MW167.01  
 2½LT V7391L Nominal concentration 0.1N (0.0167M) ±0.5% @ 20°C  
 Dgr H:350 KBrO<sub>3</sub> 2.784 g/LT  
 P:201-308+313 KBr 10 g/LT  
 Application: Reagent for volumetric titrimetry



### Potassium Bromate/Bromide 0.05N/0.1N/0.2N PrimAg-TvR

#### tri-concentrate

C6482

6x120ml C6482Q KBrO<sub>3</sub> MW 167.01  
 Dgr H:350 0.833N (0.139M) nominal concentrate ±0.5% @ 20°C  
 P:201-308+313 KBrO<sub>3</sub> 23.20 g/LT  
 KBr 83 g/LT



Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:

Dilute to 2.0LT yields 0.05N (0.0083M)

Dilute to 1.0LT yields 0.1N (0.0167M)

Dilute to 0.5LT yields 0.2N (0.0333M)

Application: Concentrate for volumetric titrimetry

*ISO 17034 accredited SI-traceable*

*Pack contains 6 vials each containing 120ml of concentrate.*

### Potassium Chloride 0.001M PrimAg-TvR

V7506

1LT V7506M KCl MW 74.55  
 2½LT V7506L Nominal concentration 0.001M (0.001N) ±0.5% @ 20°C  
 5LT V7506K KCl 0.0746 g/LT

Application: Reagent for volumetric titrimetry

*ISO 17034 accredited SI-traceable*

### Potassium Dichromate 0.025N PrimAg-TvR

V4482

1LT V4482M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> MW294.18  
 2½LT V4482L Nominal concentration 0.025N (0.004167M) ±0.5% @ 20°C  
 5LT V4482K K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> 1.226 g/LT  
 10LT V4482J Application: Reagent for volumetric titrimetry  
 Dgr H:350-340-360FD-334-317-413 *ISO 17034 accredited SI-traceable*  
 P:201-273-280g-302+352-304+341-308+313



### Potassium Dichromate 0.1N PrimAg-TvR

V5364

1LT V5364M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> MW294.18  
 2½LT V5364L Nominal concentration 0.1N (0.016667M) ±0.5% @ 20°C  
 5LT V5364K K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> 4.903 g/LT  
 10LT V5364J Application: Reagent for volumetric titrimetry  
 Dgr H:317-334-340-350-360FD-412 *ISO 17034 accredited SI-traceable*  
 P:201-273-302+352-308+313



### Potassium Dichromate 0.125N PrimAg-TvR

V6475

1LT V6475M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> MW294.18  
 2½LT V6475L Nominal concentration 0.125N (0.020833M) ±0.5% @ 20°C  
 5LT V6475K K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> 6.129 g/LT  
 10LT V6475J Application: COD analysis  
 Dgr H:317-334-340-350-360FD-412 *ISO 17034 accredited SI-traceable*  
 P:201-273-302+352-308+313



### Potassium Dichromate 0.25N PrimAg-TvR

V7786

1LT V7786M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> MW294.18  
 2½LT V7786L Nominal concentration 0.25N (0.04167M) ±0.5% @ 20°C  
 5LT V7786K K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> 12.26 g/LT  
 10LT V7786J Application: Reagent for volumetric titrimetry  
 Dgr H:315-319-317-334-340-350-360FD-373-412 *ISO 17034 accredited SI-traceable*  
 P:201-273-302+352-308+313



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Potassium Hydroxide 0.01M PrimAg-TvR

V5784

1LT V5784M  
2½LT V5784L  
5LT V5784K  
10LT V5784J

KOH MW 56.11  
Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
KOH 0.5611 g/LT  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Potassium Hydroxide 0.05M PrimAg-TvR

V6675

1LT V6675M  
2½LT V6675L  
5LT V6675K  
10LT V6675J

KOH MW 56.11  
Nominal concentration 0.05M (0.05N) ±0.5% @ 20°C  
KOH 2.806 g/LT  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Potassium Hydroxide 0.1M PrimAg-TvR

V6261

1LT V6261M  
2½LT V6261L  
5LT V6261K  
10LT V6261J  
Wing H:315-319  
P:302+352-305+351+338

KOH MW 56.11  
Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
KOH 5.611 g/LT  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*



### Potassium Hydroxide 0.5M PrimAg-TvR

V5169

1LT V5169M  
2½LT V5169L  
5LT V5169K  
10LT V5169J  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-309+310

KOH MW 56.11  
Nominal concentration 0.5M (0.5N) ±0.5% @ 20°C  
KOH 28.06 g/LT  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*



### Potassium Hydroxide 1M PrimAg-TvR

V3360

1LT V3360M  
2½LT V3360L  
5LT V3360K  
10LT V3360J  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-309+310

KOH MW 56.11  
Nominal concentration 1M (1N) ±0.5% @ 20°C  
KOH 56.11 g/LT  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*



### Potassium Hydroxide 0.5M PrimAg-TvR in industrial denatured alcohol 99

V5322

1LT V5322M  
2½LT V5322L  
Dgr H:225-290-314-302-371  
P:210-260v-280-301+330+331-305+351+338-309+310

KOH MW 56.11  
Nominal concentration 0.5M (0.5N) ±0.5% @ 20°C  
KOH 28.06 g/LT  
Solvent: industrial denatured alcohol 99  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable Not for export.*



### Potassium Hydroxide 0.1M PrimAg-TvR in methanol

V3528

1LT V3528M  
2½LT V3528L  
Dgr H:225-301+311+331-315-319-370  
P:210-280f-302+352-305+351+338-309+310-403+235

KOH MW 56.11  
Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
KOH 5.611 g/LT  
Solvent: CH<sub>3</sub>OH  
K<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Potassium Hydroxide 0.5M PrimAg-TvR

in methanol

V4326

1LT V4526M KOH MW 56.11  
 2½LT V4526L Nominal concentration 0.5M (0.5N) ±0.5% @ 20°C  
 Dgr H:225-290-314-301+311+331-370 KOH 28.06 g/LT  
 Solvent: CH3OH  
 P:210-280-302+352-305+351+338-309+310-403+235 K2CO3 (at time of manufacture) not detected  
 Protect from atmospheric CO2.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Potassium Hydroxide 1M PrimAg-TvR

in methanol

V7601

1LT V7601M KOH MW 56.11  
 2½LT V7601L Nominal concentration 1M (1N) ±0.5% @ 20°C  
 Dgr H:225-290-314-301+311+331-370 KOH 56.11 g/LT  
 Solvent: CH3OH  
 P:210-280-302+352-305+351+338-309+310-403+235 K2CO3 (at time of manufacture) not detected  
 Protect from atmospheric CO2.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Potassium Hydroxide 0.01M PrimAg-TvR

in propan-2-ol

V6850

1LT V6850M KOH MW 56.11  
 2½LT V6850L Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 Dgr H:225-319-336 KOH 0.5611 g/LT  
 P:210-233-305+351+338 Solvent: (CH3)2CHOH  
 K2CO3 (at time of manufacture) not detected  
 Protect from atmospheric CO2.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Potassium Hydroxide 0.1M PrimAg-TvR

in propan-2-ol

V5749

1LT V5749M KOH MW 56.11  
 2½LT V5749L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 Dgr H:225-315-319-336 KOH 5.611 g/LT  
 P:210-233-302+352-305+351+338 Solvent: (CH3)2CHOH  
 K2CO3 (at time of manufacture) not detected  
 Protect from atmospheric CO2.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Potassium Iodate 0.1N PrimAg-TvR

V4217

1LT V4217M KIO3 MW 214.00  
 2½LT V4217L Nominal concentration 0.1N (0.0167M) ±0.5% @ 20°C  
 5LT V4217K KIO3 3.574 g/LT  
 10LT V4217J Stabiliser: HgCl2 10 ppm  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

### Potassium Iodate 0.2N PrimAg-TvR

V5318

1LT V5318M KIO3 MW 214.00  
 2½LT V5318L Nominal concentration 0.2N (0.0333M) ±0.5% @ 20°C  
 5LT V5318K KIO3 7.126 g/LT  
 10LT V5318J Stabiliser: HgCl2 10 ppm  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

### Potassium Iodate 0.3N PrimAg-TvR

V6439

1LT V6439M KIO3 MW 214.00  
 2½LT V6439L Nominal concentration 0.3N (0.05M) ±0.5% @ 20°C  
 5LT V6439K KIO3 10.70 g/LT  
 10LT V6439J Stabiliser: HgCl2 10 ppm  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Potassium Iodate/Iodide 0.02N PrimAg-TvR

V6650

1LT V6650M KIO3 MW 214.00  
 2½LT V6650L Nominal concentration KIO3 0.02N (0.00333M) ±0.5% @ 20°C  
 KIO3 0.7126 g/LT  
 KI 7.0 g/LT  
 Stabiliser: NaHCO3 0.5 g/LT + HgCl2 10 mg/LT  
 Application: BS 1427 (1962) Determination of sulphite in boiler water  
 ISO17034 accreditedSI-traceable

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Potassium Methoxide 0.1M PrimAg-TvR in toluene/methanol 83:17

V5248

500ml V5248P  
Dgr H:225-302+312+332-304-315-  
336-361d-370  
P:210-240-280f-301+310-302+352-  
309+310-403-235

CH3OK FW 70.14  
Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
CH3OK 7.014 g/LT  
Solvent: C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>/CH<sub>3</sub>OH 83:17 v/v  
Protect from atmospheric CO<sub>2</sub>.



Application: Determination of weak carboxylic acids and their anhydrides, amides, imides and sulphonamides  
*ISO17034 accredited SI-traceable*

### Potassium Permanganate 0.02M PrimAg-TvR

V6743

1LT V6743M  
2½LT V6743L  
H:41  
2  
P:273

KMnO<sub>4</sub> MW 158.03  
Nominal concentration 0.02M (0.1N) ±0.5% @ 20°C  
KMnO<sub>4</sub> 3.161 g/LT  
Store in dark.  
Application: Reagent for volumetric titrimetry

*ISO17034 accredited SI-traceable*

### Potassium Permanganate 0.035M PrimAg-TvR

V6563

1LT V6563M  
2½LT V6563L  
H:41  
2  
P:273

KMnO<sub>4</sub> MW 158.03  
Nominal concentration 0.035M (0.175N) ±0.5% @ 20°C  
KMnO<sub>4</sub> 5.531 g/LT  
Store in dark.  
Application: Reagent for volumetric titrimetry

*ISO17034 accredited SI-traceable*

### Potassium Permanganate 0.01M/0.02M/0.04M PrimAg-TvR tri-concentrate

C7854

6x100ml C7854Q  
H:41  
1  
P:273

KMnO<sub>4</sub> MW 158.03  
0.2M (1N) nominal concentrate ±0.5% @ 20°C  
KMnO<sub>4</sub> 31.61 g/LT



Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:

Dilute to 2.0LT yields 0.01M (0.05N)  
Dilute to 1.0LT yields 0.02M (0.1N)  
Dilute to 0.5LT yields 0.04M (0.2N)

Store in dark.

Application: Concentrate for volumetric titrimetry

*ISO 17034 accredited SI-traceable*

Pack contains 6 vials each containing 100ml of concentrate.

### Potassium Permanganate 0.2M PrimAg-TvR

V5834

1LT V5834M  
2½LT V5834L  
H:41  
1  
P:273

KMnO<sub>4</sub> MW 158.03  
Nominal concentration 0.2M (1N) ±0.5% @ 20°C  
KMnO<sub>4</sub> 31.61 g/LT  
Store in dark.  
Application: Reagent for volumetric titrimetry



*ISO17034 accredited SI-traceable*

### Potassium Thiocyanate 0.05M PrimAg-TvR

V6824

1LT V6824M  
2½LT V6824L  
5LT V6824K  
10LT V6824J

KSCN MW 97.18  
Nominal concentration 0.05M (0.05N) ±0.5% @ 20°C  
KSCN 4.859 g/LT  
Application: Reagent for volumetric titrimetry

*ISO17034 accredited SI-traceable*

### Potassium Thiocyanate 0.1M PrimAg-TvR

V7835

1LT V7835M  
2½LT V7835L  
5LT V7835K  
10LT V7835J

KSCN MW 97.18  
Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
KSCN 9.718 g/LT  
Application: Reagent for volumetric titrimetry

*ISO17034 accredited SI-traceable*

Sodium Dodecylbenzenesulphonate \*M (see Dodecylbenzenesulphonic Acid sodium salt \*M)

Sodium Lauryl Sulphate \*M (see Sodium Dodecyl Sulphate \*M)



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Silver Nitrate 0.001M PrimAg-TvR in propan-2-ol/water 90:10

V6529

1LT V6529M AgNO<sub>3</sub> MW 169.87  
 2½LT V6529L Nominal concentration 0.001M (0.001N) ±0.5% @ 20°C  
 Dgr H:225-319-336 AgNO<sub>3</sub> 0.1699 g/LT  
 P:210-233-305+351+338 Solvent: (CH<sub>3</sub>)<sub>2</sub>CHOH/H<sub>2</sub>O 90:10 v/v  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable



### Silver Nitrate 0.01M PrimAg-TvR

V5476

1LT V5476M AgNO<sub>3</sub> MW169.87  
 2½LT V5476L Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 10LT V5476J AgNO<sub>3</sub> 1.699 g/LT  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Silver Nitrate 0.01M PrimAg-TvR in propan-2-ol/water 90:10

V7436

1LT V7436M AgNO<sub>3</sub> MW 169.87  
 2½LT V7436L Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 Dgr H:225-315-319-336-410 AgNO<sub>3</sub> 1.699 g/LT  
 P:210-233-273-302+352-305+351+338 Solvent: (CH<sub>3</sub>)<sub>2</sub>CHOH/H<sub>2</sub>O 90:10 v/v  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable



### Silver Nitrate 0.02M PrimAg-TvR

V4018

1LT V4018M AgNO<sub>3</sub> MW169.87  
 2½LT V4018L Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 10LT V4018J AgNO<sub>3</sub> 3.397 g/LT  
 Wng H:315-319-411 Application: Reagent for volumetric titrimetry  
 P:273-302+352-305+351+338 ISO17034 accredited SI-traceable



### Silver Nitrate 0.0282M PrimAg-TvR 1 ml = 1 mg Cl<sup>-</sup>

V5331

1LT V5331M AgNO<sub>3</sub> MW169.87  
 2½LT V5331L Nominal concentration 0.0282M (0.0282N) ±0.5% @ 20°C  
 10LT V5331J AgNO<sub>3</sub> 4.790 g/LT  
 Wng H:315-319-411 Application: Reagent for volumetric titrimetry  
 P:273-302+352-305+351+338 ISO17034 accredited SI-traceable



### Silver Nitrate 0.05M PrimAg-TvR

V3674

1LT V3674M AgNO<sub>3</sub> MW169.87  
 2½LT V3674L Nominal concentration 0.05M (0.05N) ±0.5% @ 20°C  
 10LT V3674J AgNO<sub>3</sub> 8.494 g/LT  
 Wng H:315-319-411 Application: Reagent for volumetric titrimetry  
 P:273-302+352-305+351+338 ISO17034 accredited SI-traceable



### Silver Nitrate 0.1M PrimAg-TvR

V6666

1LT V6666M AgNO<sub>3</sub> MW169.87  
 2½LT V6666L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 10LT V6666J AgNO<sub>3</sub> 16.99 g/LT  
 Wng H:315-319-411 Application: Reagent for volumetric titrimetry  
 P:273-302+352-305+351+338 ISO17034 accredited SI-traceable



### Silver Nitrate 0.05M/0.1M/0.2M PrimAg-TvR

C7777

6x100ml C7777Q **tri-concentrate**  
 AgNO<sub>3</sub> MW 169.87  
 Dgr H:314-410 1M (1N) nominal concentrate ±0.5% @ 20°C  
 P:273-280c-301+330+331-305+351+338-309+310 AgNO<sub>3</sub> 169.9 g/LT  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 2.0LT yields 0.05M (0.05N)  
 Dilute to 1.0LT yields 0.1M (0.1N)  
 Dilute to 0.5LT yields 0.2M (0.2N)  
 Application: Concentrate for volumetric titrimetry  
 ISO 17034 accredited SI-traceable  
 Pack contains 6 vials each containing 100ml of concentrate.



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Silver Nitrate 0.25M PrimAg-TvR

V6785

1LT V6785M AgNO<sub>3</sub> MW169.87  
 2½LT V6785L Nominal concentration 0.25M (0.25N) ±0.5% @ 20°C  
 Wng H:315-319-410 AgNO<sub>3</sub> 42.47 g/LT  
 P:273-302+352-305+351+338 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable



### Silver Nitrate 1M PrimAg-TvR

V6752

500ml V6752P AgNO<sub>3</sub> MW169.87  
 1LT V6752M Nominal concentration 1M (1N) ±0.5% @ 20°C  
 2½LT V6752L AgNO<sub>3</sub> 169.9 g/LT  
 Dgr H:314-410 Application: Reagent for volumetric titrimetry  
 P:273-280c-301+330+331-305+351+338-309+310 ISO17034 accredited SI-traceable



### Sodium Arsenite 0.0125M PrimAg-TvR

V6537

1LT V6537M NaAsO<sub>2</sub> MW129.91  
 2½LT V6537L Nominal concentration 0.0125M (0.025N) ±0.5% @ 20°C  
 5LT V6537K NaAsO<sub>2</sub> 1.624 g/LT  
 10LT V6537J Application: Reagent for volumetric titrimetry  
 Dgr H:350 ISO17034 accredited SI-traceable  
 P:201-308+313



### Sodium Arsenite 0.025M PrimAg-TvR

V5158

1LT V5158M NaAsO<sub>2</sub> MW129.91  
 2½LT V5158L Nominal concentration 0.025M (0.05N) ±0.5% @ 20°C  
 5LT V5158K NaAsO<sub>2</sub> 3.248 g/LT  
 10LT V5158J Application: Reagent for volumetric titrimetry  
 Dgr H:350-412 ISO17034 accredited SI-traceable  
 P:201-273-308+313



### Sodium Arsenite 0.05M PrimAg-TvR

V3780

1LT V3780M NaAsO<sub>2</sub> MW129.91  
 2½LT V3780L Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C  
 5LT V3780K NaAsO<sub>2</sub> 6.496 g/LT  
 10LT V3780J Application: Reagent for volumetric titrimetry  
 Dgr H:350-412 ISO17034 accredited SI-traceable  
 P:201-273-308+313



### Sodium Carbonate 0.02M PrimAg-TvR

V5367

1LT V5367M Na<sub>2</sub>CO<sub>3</sub> MW105.99  
 2½LT V5367L Nominal concentration 0.02M (0.04N) ±0.5% @ 20°C  
 5LT V5367K Na<sub>2</sub>CO<sub>3</sub> 2.120 g/LT  
 10LT V5367J Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Sodium Carbonate 0.5M PrimAg-TvR

V4500

1LT V4500M Na<sub>2</sub>CO<sub>3</sub> MW105.99  
 2½LT V4500L Nominal concentration 0.5M (1N) ±0.5% @ 20°C  
 5LT V4500K Na<sub>2</sub>CO<sub>3</sub> 53.00 g/LT  
 10LT V4500J Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Sodium Chloride 0.01M PrimAg-TvR

V0456

1LT V0456M NaCl MW 58.44  
 Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 NaCl 0.5844 g/LT  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accredited SI-traceable

### Sodium Chloride 0.02M PrimAg-TvR

V2467

1LT V2467M NaCl MW58.44  
 Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 NaCl 1.169 g/LT  
 Application: Reagent for volumetric titrimetry  
 ISO 17034 accredited SI-traceable

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sodium Chloride 0.1M PrimAg-TvR

V6453

1LT V6453M NaCl MW 58.44  
 Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 NaCl 5.844 g/LT  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Sodium Chloride 1M PrimAg-TvR

V3445

1LT V3445M NaCl MW 58.44  
 Nominal concentration 1M (1N) ±0.5% @ 20°C  
 NaCl 58.44 g/LT  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Sodium Dodecyl Sulphate 0.004M PrimAg-TvR

V7104

1LT V7104M **(Sodium Lauryl Sulphate 0.004M)**  
 2½LT V7104L CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na MW 288.38  
 Nominal concentration 0.004M ±0.5% @ 20°C  
 CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na 1.154 g/LT  
 Store at 15-25°C.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*  
*Product may precipitate at low temperatures. Gentle warming of the container will redissolve the solid and will not affect the certified concentration.*

### Sodium Dodecyl Sulphate 0.05M PrimAg-TvR

V5306

1LT V5306M **(Sodium Lauryl Sulphate 0.05M)**  
 2½LT V5306L CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na MW 288.38  
 Nominal concentration 0.05M ±0.5% @ 20°C  
 CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na 14.42 g/LT  
 Store at 15-25°C.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*  
*Product may precipitate at low temperatures. Gentle warming of the container will redissolve the solid and will not affect the certified concentration.*

### Sodium Dodecyl Sulphate 0.1M PrimAg-TvR

V6425

1LT V6425M **(Sodium Lauryl Sulphate 0.1M)**  
 2½LT V6425L CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na MW 288.38  
 Nominal concentration 0.1M ±0.5% @ 20°C  
 CH<sub>3</sub>(CH<sub>2</sub>)<sub>11</sub>OSO<sub>3</sub>.Na 28.84 g/LT  
 Store at 15-25°C.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*  
*Product may precipitate at low temperatures. Gentle warming of the container will redissolve the solid and will not affect the certified concentration.*

### Sodium Hydroxide 0.01M PrimAg-TvR

V5902

1LT V5902M NaOH MW 40.00  
 2½LT V5902L Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 5LT V5902K NaOH 0.4000 g/LT  
 10LT V5902J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Sodium Hydroxide 0.02 M PrimAg-TvR

V6136

1LT V6136M NaOH MW 40.00  
 2½LT V6136L Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 5LT V6136K NaOH 0.8000 g/LT  
 10LT V6136J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

### Sodium Hydroxide 0.04M PrimAg-TvR high accuracy

V9404

1LT V9404M NaOH MW 40.00  
 2½LT V9404L Nominal concentration 0.04M (0.04N) ±0.01% @ 20°C  
 NaOH 1.6000 g/LT  
 Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
 To maintain high accuracy protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sodium Hydroxide 0.05M PrimAg-TvR

V3573

1LT V3573M NaOH MW 40.00  
 2½LT V3573L Nominal concentration 0.05M (0.05N) ±0.5% @ 20°C  
 5LT V3573K NaOH 2.000 g/LT  
 10LT V3573J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Sodium Hydroxide 0.1M PrimAg-TvR

V6701

1LT V6701M NaOH MW 40.00  
 2½LT V6701L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 5LT V6701K NaOH 4.000 g/LT  
 10LT V6701J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Sodium Hydroxide 0.05M/0.1M PrimAg-TvR di-concentrate

C7812

6x100ml C7812Q NaOH MW 40.00  
 Dgr H:290-314 1M (1N) nominal concentrate ±0.5% @ 20°C  
 P:280c-301+330+331-305+351+338- NaOH 40.00 g/LT  
 310 Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
 Protect from atmospheric CO<sub>2</sub>.  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 2.0LT yields 0.05M (0.05N)  
 Dilute to 1.0LT yields 0.1M (0.1N)  
 Application: Concentrate for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*  
 Pack contains 6 vials each containing 100ml of concentrate.

### Sodium Hydroxide 0.111M PrimAg-TvR 1 ml = 1 mg lactic acid

V4153

1LT V4153M NaOH MW 40.00  
 2½LT V4153L Nominal concentration 0.111M (0.111N) ±0.5% @ 20°C  
 5LT V4153K NaOH 4.440 g/LT  
 10LT V4153J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Determination of acidity of milk products  
*ISO 17034 accredited SI-traceable*

### Sodium Hydroxide 0.125M PrimAg-TvR

V5428

1LT V5428M NaOH MW 40.00  
 2½LT V5428L Nominal concentration 0.125M (0.125N) ±0.5% @ 20°C  
 5LT V5428K NaOH 5.000 g/LT  
 10LT V5428J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Sodium Hydroxide 0.2M PrimAg-TvR

V6384

1LT V6384M NaOH MW 40.00  
 2½LT V6384L Nominal concentration 0.2M (0.2N) ±0.5% @ 20°C  
 5LT V6384K NaOH 8.000 g/LT  
 10LT V6384J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Sodium Hydroxide 0.2M PrimAg-TvR high accuracy

V9420

1LT V9420M NaOH MW 40.00  
 2½LT V9420L Nominal concentration 0.2M (0.2N) ±0.01% @ 20°C  
 Wng H:290-315-319 NaOH 8.0000 g/LT  
 P:302+352-305+351+338 Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
 To maintain high accuracy protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

### Sodium Hydroxide 0.25M PrimAg-TvR

V5217

1LT V5217M NaOH MW 40.00  
 2½LT V5217L Nominal concentration 0.25M (0.25N) ±0.5% @ 20°C  
 5LT V5217K NaOH 10.00 g/LT  
 Wng H:290-315-319 Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sodium Hydroxide 0.313M PrimAg-TvR

V6061

1LT V6061M  
2½LT V6061L  
5LT V6061K  
10LT V6061J  
Wng H:290-315-319  
P:302+352-305+351+338

NaOH MW 40.00  
Nominal concentration 0.313M (0.313N) ±0.5% @ 20°C  
NaOH 12.52 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Sodium Hydroxide 0.33M PrimAg-TvR

V4537

1LT V4537M  
2½LT V4537L  
5LT V4537K  
10LT V4537J  
Wng H:290-315-319  
P:302+352-305+351+338

NaOH MW 40.00  
Nominal concentration 0.33M (0.33N) ±0.5% @ 20°C  
NaOH 13.20 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Sodium Hydroxide 0.5M PrimAg-TvR

V3858

1LT V3858M  
2½LT V3858L  
5LT V3858K  
10LT V3858J  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-310

NaOH MW 40.00  
Nominal concentration 0.5M (0.5N) ±0.5% @ 20°C  
NaOH 20.00 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Sodium Hydroxide 1M PrimAg-TvR

V6713

1LT V6713M  
2½LT V6713L  
5LT V6713K  
10LT V6713J  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-310

NaOH MW 40.00  
Nominal concentration 1M (1N) ±0.5% @ 20°C  
NaOH 40.00 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Sodium Hydroxide 0.5M/1M PrimAg-TvR di-concentrate

C7825

6x100ml C7825Q  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-310

NaOH MW 40.00  
10M (10N) nominal concentrate ±0.5% @ 20°C  
NaOH 400.0 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
Dilute to 2.0LT yields 0.5M (0.5N)  
Dilute to 1.0LT yields 1M (1N)  
Application: Concentrate for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*  
Pack contains 6 vials each containing 100ml of concentrate.



### Sodium Hydroxide 2M PrimAg-TvR

V5308

1LT V5308M  
2½LT V5308L  
5LT V5308K  
10LT V5308J  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-310

NaOH MW 40.00  
Nominal concentration 2M (2N) ±0.5% @ 20°C  
NaOH 80.00 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



### Sodium Hydroxide 4M PrimAg-TvR

V5357

1LT V5357M  
2½LT V5357L  
5LT V5357K  
10LT V5357J  
Dgr H:290-314  
P:280c-301+330+331-305+351+338-310

NaOH MW 40.00  
Nominal concentration 4M (4N) ±0.5% @ 20°C  
NaOH 160.0 g/LT  
Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
Protect from atmospheric CO<sub>2</sub>.  
Application: Reagent for volumetric titrimetry



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sodium Hydroxide 5M PrimAg-TvR

V5576

1LT V5576M NaOH MW 40.00  
 2½LT V5576L Nominal concentration 5M (5N) ±0.5% @ 20°C  
 5LT V5576K NaOH 200.0 g/LT  
 10LT V5576J Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
 Dgr H:290-314 Protect from atmospheric CO<sub>2</sub>.  
 P:280c-301+330+331-305+351+338-310 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Sodium Hydroxide 9M PrimAg-TvR

V4734

1LT V4734M NaOH MW 40.00  
 2½LT V4734L Nominal concentration 9M (9N) ±0.5% @ 20°C  
 5LT V4734K NaOH 360.0 g/LT  
 Dgr H:290-314 Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
 P:280c-301+330+331-305+351+338-310 Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Sodium Hydroxide 10M PrimAg-TvR

V6326

1LT V6326M NaOH MW 40.00  
 2½LT V6326L Nominal concentration 10M (10N) ±0.5% @ 20°C  
 5LT V6326K NaOH 400.0 g/LT  
 Dgr H:290-314 Na<sub>2</sub>CO<sub>3</sub> (at time of manufacture) not detected  
 P:280c-301+330+331-305+351+338-310 Protect from atmospheric CO<sub>2</sub>.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable



### Sodium Thiosulphate 0.005M PrimAg-TvR

V2473

1LT V2473M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V2473L Nominal concentration 0.005M (0.005N) ±0.5% @ 20°C  
 5LT V2473K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 0.7906 g/LT  
 Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

### Sodium Thiosulphate 0.01M PrimAg-TvR

V6360

1LT V6360M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V6360L Nominal concentration 0.01M (0.01N) ±0.5% @ 20°C  
 5LT V6360K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 1.581 g/LT  
 10LT V6360J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

### Sodium Thiosulphate 0.02M PrimAg-TvR

V1375

1LT V1375M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V1375L Nominal concentration 0.02M (0.02N) ±0.5% @ 20°C  
 5LT V1375K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 3.162 g/LT  
 10LT V1375J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

### Sodium Thiosulphate 0.025M PrimAg-TvR

V3682

1LT V3682M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V3682L Nominal concentration 0.025M (0.025N) ±0.5% @ 20°C  
 5LT V3682K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 3.953 g/LT  
 10LT V3682J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

### Sodium Thiosulphate 0.05M PrimAg-TvR

V5677

1LT V5677M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V5677L Nominal concentration 0.05M (0.05N) ±0.5% @ 20°C  
 5LT V5677K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 7.906 g/LT  
 10LT V5677J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
 ISO17034 accreditedSI-traceable

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sodium Thiosulphate 0.1M PrimAg-TvR

V7322

1LT V7322M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V7322L Nominal concentration 0.1M (0.1N) ±0.5% @ 20°C  
 5LT V7322K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 15.81 g/LT  
 10LT V7322J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*

### Sodium Thiosulphate 0.05M/0.1M/0.2M PrimAg-TvR

tri-concentrate

C8433

6x100ml C8433Q Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 1M (1N) nominal concentration ±0.5% @ 20°C  
 Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 158.1 g/LT  
 Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 2.0LT yields 0.05M (0.05N)  
 Dilute to 1.0LT yields 0.1M (0.1N)  
 Dilute to 0.5LT yields 0.2M (0.2N)  
 Application: Concentrate for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*  
 Pack contains 6 vials each containing 100ml of concentrate.

### Sodium Thiosulphate 0.2M PrimAg-TvR

V4421

1LT V4421M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V4421L Nominal concentration 0.2M (0.2N) ±0.5% @ 20°C  
 5LT V4421K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 31.62 g/LT  
 10LT V4421J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*

### Sodium Thiosulphate 0.125M/0.25M/0.5M PrimAg-TvR

tri-concentrate

C5534

6x100ml C5534Q Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2.5M (2.5N) nominal concentration ±0.5% @ 20°C  
 Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 395.3 g/LT  
 Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Dilution of the vial contents with distilled water using class A glassware gives the following working strengths:  
 Dilute to 2.0LT yields 0.125M (0.125N)  
 Dilute to 1.0LT yields 0.25M (0.25N)  
 Dilute to 0.5LT yields 0.5M (0.5N)  
 Application: Concentrate for volumetric titrimetry  
*ISO 17034 accredited SI-traceable*  
 Pack contains 6 vials each containing 100ml of concentrate.

### Sodium Thiosulphate 1M PrimAg-TvR

V8024

1LT V8024M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> MW 158.11  
 2½LT V8024L Nominal concentration 1M (1N) ±0.5% @ 20°C  
 5LT V8024K Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> 158.1 g/LT  
 10LT V8024J Stabiliser: Na<sub>2</sub>CO<sub>3</sub> 0.2 g/LT  
 Protect from atmospheric microbial contamination.  
 Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*

### Sulphuric Acid 0.005M PrimAg-TvR

V8443

1LT V8443M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8443L Nominal concentration 0.005M (0.01N) ±0.5% @ 20°C  
 5LT V8443K H<sub>2</sub>SO<sub>4</sub> 0.4904 g/LT  
 10LT V8443J Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*

### Sulphuric Acid 0.01M PrimAg-TvR

V8554

1LT V8554M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8554L Nominal concentration 0.01M (0.02N) ±0.5% @ 20°C  
 5LT V8554K H<sub>2</sub>SO<sub>4</sub> 0.9807 g/LT  
 10LT V8554J Application: Reagent for volumetric titrimetry  
*ISO17034 accredited SI-traceable*



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sulphuric Acid 0.05M PrimAg-TvR

V8639

1LT V8639M	H2SO4 MW98.07
2½LT V8639L	Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C
5LT V8639K	H2SO4 4.904 g/LT
10LT V8639J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>

### Sulphuric Acid 0.1M PrimAg-TvR

V8641

1LT V8641M	H2SO4 MW98.07
2½LT V8641L	Nominal concentration 0.1M (0.2N) ±0.5% @ 20°C
5LT V8641K	H2SO4 9.807 g/LT
10LT V8641J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>

### Sulphuric Acid 0.05M/0.1M/0.2M PrimAg-TvR

tri-concentrate

C8752

6x100ml C8752Q	H2SO4 MW 98.07
Wng H:290-315-319	1M (2N) nominal concentrate ±0.5% @ 20°C
P:302+352-305+351+338	H2SO4 98.07 g/LT



Dilution of the vial contents with distilled water using class A glassware gives

the following working strengths:

Dilute to 2.0LT yields 0.05M (0.1N)

Dilute to 1.0LT yields 0.1M (0.2N)

Dilute to 0.5LT yields 0.2M (0.4N)

Application: Concentrate for volumetric titrimetry

*ISO 17034 accredited SI-traceable*

Pack contains 6 vials each containing 100ml of concentrate.

### Sulphuric Acid 0.125M PrimAg-TvR

V7530

1LT V7530M	H2SO4 MW98.07
2½LT V7530L	Nominal concentration 0.125M (0.25N) ±0.5% @ 20°C
5LT V7530K	H2SO4 12.26 g/LT
10LT V7530J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>

### Sulphuric Acid 0.1275M PrimAg-TvR

V2386

1LT V2386M	H2SO4 MW98.07
2½LT V2386L	Nominal concentration 0.1275M (0.255N) ±0.5% @ 20°C
5LT V2386K	H2SO4 12.50 g/LT
10LT V2386J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>

### Sulphuric Acid 0.25M PrimAg-TvR

V6812

1LT V6812M	H2SO4 MW 98.07
2½LT V6812L	Nominal concentration 0.25M (0.5N) ±0.5% @ 20°C
5LT V6812K	H2SO4 24.52 g/LT
10LT V6812J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>

### Sulphuric Acid 0.5M PrimAg-TvR

V5345

1LT V5345M	H2SO4 MW 98.07
2½LT V5345L	Nominal concentration 0.5M (1N) ±0.5% @ 20°C
5LT V5345K	H2SO4 49.04 g/LT
10LT V5345J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>

### Sulphuric Acid 0.25M/0.5M/1M PrimAg-TvR

tri-concentrate

C5456

6x100ml C5456Q	H2SO4 MW 98.07
Dgr H:290-314	5M (10N) nominal concentrate ±0.5% @ 20°C
P:280c-301+330+331-305+351+338-309+310	H2SO4 490.4 g/LT



Dilution of the vial contents with distilled water using class A glassware gives

the following working strengths:

Dilute to 2.0LT yields 0.25M (0.5N)

Dilute to 1.0LT yields 0.5M (1N)

Dilute to 0.5LT yields 1M (2N)

Application: Concentrate for volumetric titrimetry

*ISO 17034 accredited SI-traceable*

Pack contains 6 vials each containing 100ml of concentrate.

### Sulphuric Acid 1M PrimAg-TvR

V8700

1LT V8700M	H2SO4 MW 98.07
2½LT V8700L	Nominal concentration 1M (2N) ±0.5% @ 20°C
5LT V8700K	H2SO4 98.07 g/LT
10LT V8700J	Application: Reagent for volumetric titrimetry <i>ISO17034 accredited SI-traceable</i>
Wng H:290-315-319	
P:302+352-305+351+338	



## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sulphuric Acid 2M PrimAg-TvR

V8115

1LT V8115M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8115L Nominal concentration 2M (4N) ±0.5% @ 20°C  
 5LT V8115K H<sub>2</sub>SO<sub>4</sub> 196.1 g/LT  
 10LT V8115J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 2.5M PrimAg-TvR

V8236

1LT V8236M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8236L Nominal concentration 2.5M (5N) ±0.5% @ 20°C  
 5LT V8236K H<sub>2</sub>SO<sub>4</sub> 245.2 g/LT  
 10LT V8236J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 3M PrimAg-TvR

V7455

1LT V7455M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V7455L Nominal concentration 3M (6N) ±0.5% @ 20°C  
 5LT V7455K H<sub>2</sub>SO<sub>4</sub> 294.2 g/LT  
 10LT V7455J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 4M PrimAg-TvR

V8665

1LT V8665M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8665L Nominal concentration 4M (8N) ±0.5% @ 20°C  
 5LT V8665K H<sub>2</sub>SO<sub>4</sub> 392.3 g/LT  
 10LT V8665J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 5M PrimAg-TvR

V6546

1LT V6546M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V6546L Nominal concentration 5M (10N) ±0.5% @ 20°C  
 5LT V6546K H<sub>2</sub>SO<sub>4</sub> 490.4 g/LT  
 10LT V6546J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 12M PrimAg-TvR

V8912

1LT V8912M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8912L Nominal concentration 12M (24N) ±0.5% @ 20°C  
 5LT V8912K H<sub>2</sub>SO<sub>4</sub> 1177 g/LT  
 10LT V8912J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Zinc Acetate 0.05M PrimAg-TvR

V6127

1LT V6127M (CH<sub>3</sub>COO)<sub>2</sub>Zn MW183.47  
 2½LT V6127L Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C  
 5LT V6127K (CH<sub>3</sub>COO)<sub>2</sub>Zn 9.174 g/LT  
 10LT V6127J Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

## ROMIL PrimAg®-TvR Traceable Volumetric Reagents Specifications

### Sulphuric Acid 2M PrimAg-TvR

V8115

1LT V8115M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8115L Nominal concentration 2M (4N) ±0.5% @ 20°C  
 5LT V8115K H<sub>2</sub>SO<sub>4</sub> 196.1 g/LT  
 10LT V8115J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 2.5M PrimAg-TvR

V8236

1LT V8236M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8236L Nominal concentration 2.5M (5N) ±0.5% @ 20°C  
 5LT V8236K H<sub>2</sub>SO<sub>4</sub> 245.2 g/LT  
 10LT V8236J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 3M PrimAg-TvR

V7455

1LT V7455M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V7455L Nominal concentration 3M (6N) ±0.5% @ 20°C  
 5LT V7455K H<sub>2</sub>SO<sub>4</sub> 294.2 g/LT  
 10LT V7455J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 4M PrimAg-TvR

V8665

1LT V8665M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8665L Nominal concentration 4M (8N) ±0.5% @ 20°C  
 5LT V8665K H<sub>2</sub>SO<sub>4</sub> 392.3 g/LT  
 10LT V8665J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 5M PrimAg-TvR

V6546

1LT V6546M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V6546L Nominal concentration 5M (10N) ±0.5% @ 20°C  
 5LT V6546K H<sub>2</sub>SO<sub>4</sub> 490.4 g/LT  
 10LT V6546J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Sulphuric Acid 12M PrimAg-TvR

V8912

1LT V8912M H<sub>2</sub>SO<sub>4</sub> MW 98.07  
 2½LT V8912L Nominal concentration 12M (24N) ±0.5% @ 20°C  
 5LT V8912K H<sub>2</sub>SO<sub>4</sub> 1177 g/LT  
 10LT V8912J Application: Reagent for volumetric titrimetry  
 Dgr H:290-314 *ISO17034 accreditedSI-traceable*  
 P:280c-301+330+331-305+351+338-309+310



### Zinc Acetate 0.05M PrimAg-TvR

V6127

1LT V6127M (CH<sub>3</sub>COO)<sub>2</sub>Zn MW183.47  
 2½LT V6127L Nominal concentration 0.05M (0.1N) ±0.5% @ 20°C  
 5LT V6127K (CH<sub>3</sub>COO)<sub>2</sub>Zn 9.174 g/LT  
 10LT V6127J Application: Reagent for volumetric titrimetry  
*ISO17034 accreditedSI-traceable*

## ROMIL Technical Grade Solvents and Acids Specifications

Acetic Acid glacial (see Acetic Acid)

### Acetic Acid tech

RS014

2½LT RS014L  
25LT RS014G  
Dgr H:226-314  
P:280c-301+330+331-305+351+338-307+310



(Acetic Acid glacial)

CH<sub>3</sub>COOH MW60.05 FP 16.7°C BP 117.9°C d 1.05 CAS [64-19-7] Assay >99% Water <0.5%

### Acetone tech

RS031

2½LT RS031L  
5LT RS031K  
10LT RS031J  
25LT RS031G  
Dgr H:225-319-336-EUH066  
P:210-233-305+351+338



(Propanone)

(CH<sub>3</sub>)<sub>2</sub>COMW 58.08 BP 56.1°C d 0.79 CAS [67-64-1] Assay >99% Water <0.8%

### Acetonitrile tech

RS046

2½LT RS046L  
25LT RS046G  
Dgr H:225-302+312+332-319  
P:210-240-302+352-305+351+338-403+233



(Methyl Cyanide)

CH<sub>3</sub>CNMW 41.05 BP 81.6°C d 0.78 CAS [75-05-8] Assay >99% Water <0.3%

### iso-Amyl Alcohol tech

RS062

2½LT RS062L  
25LT RS062G  
Wng H:226-332-335-EUH066  
P:210-304+340



(iso-Pentanol)

(CH<sub>3</sub>)CH(CH<sub>2</sub>)<sub>2</sub>OH MW 88.15 d 0.81 Water <0.8% Assay (3- and 2-methyl isomers) >97%

n-Butanol (see Butan-1-ol)

2-Butanone (see Methyl Ethyl Ketone)

n-Butyl Alcohol (see Butan-1-ol)

n-Butyl Chloride (see 1-Chlorobutane)

### Butan-1-ol tech

RS083

2½LT RS083L 25LT RS083G  
Dgr H:226-302-315-318-335-336  
P:210-280f-302+352-304+340-305+351+338-313



(n-Butanol, n-Butyl Alcohol)

CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>OHMW 74.12 BP 117.7°C d 0.81 CAS [71-36-3] Assay >99% Water <0.5%

### 1-Chlorobutane tech

RS118

2½LT RS118L  
25LT RS118G  
Dgr H:225  
P:210



(n-Butyl Chloride)

CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>Cl MW 92.57 BP 78.4°C d 0.88 CAS [109-69-3] Assay >98% Water <0.5%

### Chloroform tech

RS135

2½LT RS135L  
Dgr H:351-361d-331-302-372-319-315  
P:261v-280f-304+340-305+351+338-308+313



(Trichloromethane)

CHCl<sub>3</sub> MW119.38 BP 61.2°C d 1.48 CAS [67-66-3] Assay >98%\* Water <0.2%

\*ex stabiliser

Stabiliser: Ethanol ca. 1% w/w

Stabiliser should only be removed immediately before use by adsorption onto activated alumina.

### Cyclohexane tech

RS156

2½LT RS156L 25LT RS156G  
Dgr H:225-304-315-336-410  
P:210-233-240-273-301+310-302+352-331-403+235



C<sub>6</sub>H<sub>12</sub> MW 84.16 FP6.5°C BP 80.7°C d 0.78 CAS [110-82-7]

Assay >99% Water <0.2%

## ROMIL Technical Grade Solvents and Acids Specifications

### Dichloromethane tech

RS202

2½LT RS202L  
25LT RS202G  
Wng H:351  
P:281-308+313

**(Methylene Dichloride)**

CH<sub>2</sub>Cl<sub>2</sub>MW 84.93BP 39.6°C d 1.33 CAS [75-09-2]  
Assay >99%\* Water <0.2%  
\*ex stabiliser  
Stabiliser: Amylene ca. 50 ppm



### Diethyl Ether tech

RS220

2½LT RS220L  
25LT RS220G  
Dgr H:224-302-336-EUH019-EUH066  
P:210-240-403+235

(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>OMW 74.12 BP34.4°C d 0.71 CAS [60-29-7]  
Assay >99%\* Water <0.2%  
\*ex stabiliser  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 5 ppm



### Dimethylformamide tech

RS253

2½LT RS253L  
25LT RS253G  
Dgr H:360D-226-312+332-319  
P:201-210-302+352-305+351+338-308+313

HCON(CH<sub>3</sub>)<sub>2</sub> MW 73.09BP153.0°Cd0.95 CAS [68-12-2]  
Assay >99% Water <0.3%



### 1,4-Dioxan tech

RS297

2½LT RS297L  
25LT RS297G  
Dgr H:225-350-319-335-EUH019-EUH066  
P:210-281-305+351+338-308+313

C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>MW 88.11 FP11.8°C BP 101.3°C d 1.03 CAS [123-91-1]  
Assay >99% Water <0.2%  
Unstabilised



### Di-iso-propyl Ether tech

RS236

2½LT RS236L  
25LT RS236G  
Dgr H:225-336-EUH019-EUH066  
P:210-240-403+235

[(CH<sub>3</sub>)<sub>2</sub>CH]<sub>2</sub>O MW 102.18BP68.5°Cd0.73 CAS [108-20-3]  
Assay >98%\* Water <0.5%  
\*ex stabiliser  
Stabiliser: Butylated hydroxytoluene (BHT) ca. 5 ppm



### Ethanol absolute tech

2½LT RS314L  
25LT RS314G  
Dgr H:225  
P:210-233-240-403+235

**(Ethyl Alcohol)**

C<sub>2</sub>H<sub>5</sub>OHMW 46.07 BP 78.3°C d 0.79 CAS [64-17-5]  
Assay >99% Water <0.8%



### n-Heptane 95% tech

RS367

2½LT RS367L  
25LT RS367G  
Dgr H:225-304-315-336-410  
P:210-273-301+310-331-302+352-304+340-403+235

CH<sub>3</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub> MW 100.21BP94-98°C d 0.68 CAS [142-82-5]  
Water <0.2%  
Assay (n-isomer) ca. 95%  
Assay (all isomers) >99%



### Hexane tech

RS390

2½LT RS390L  
25LT RS390G  
Dgr H:225-304-361f-373-315-336-411  
P:210-240-273-301+310-331-302+352-403+235

C<sub>6</sub>H<sub>14</sub> BP 65-70°C d 0.66 CAS [73513-42-5]  
Water <0.2%

Comprises ca. 50% n-isomer, the remainder being predominantly other isomers of hexane.



### n-Hexane 95% tech

RS389

2½LT RS389L  
25LT RS389G  
Dgr H:225-304-361f-373-315-336-411  
P:210-240-273-301+310-331-302+352-403+235

CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub> MW 86.18BP67-70°C d 0.66 CAS [110-54-3]  
Water <0.2%

Assay (n-isomer) ca. 95%  
Assay (all isomers) >99%



## ROMIL Technical Grade Solvents and Acids Specifications

### Hydrochloric Acid 1.18 tech

RA396

2½LT RA396L HCIMW 36.46d1.18CAS[7647-01-0]  
 25LT RA396G Assay ca. 36%  
 Dgr H:290-314-335  
 P:280c-301+330+331-305+351+338-309+310



Methyl Alcohol (see Methanol)

Methyl Cyanide (see Acetonitrile)

Methylene Dichloride (see Dichloromethane)

### Methanol tech

RS409

2½LT RS409L (Methyl Alcohol)  
 25LT RS409G CH<sub>3</sub>OHMW 32.04 BP 64.5°C d 0.79 CAS [67-56-1]  
 Dgr H:225-301+311+331-370 Assay >99% Water <0.5%  
 P:210-280f-302+352-309+310-403+235



### Methyl Ethyl Ketone tech

RS493

2½LT RS493L (2-Butanone)  
 25LT RS493G CH<sub>3</sub>CH<sub>2</sub>COCH<sub>3</sub> MW 72.11 BP 79.6°C d 0.80 CAS [78-93-3]  
 Dgr H:225-319-336-EUH066 Assay >99% Water <0.3%  
 P:210-305+351+338-403+233



### Nitric Acid 1.42 tech

RA566

2½LT RA566L HNO<sub>3</sub> MW63.01d1.42CAS[7697-37-2]  
 Dgr H:272-290-331-314-EUH071 Assay ca. 69%  
 P:260c-280c-301+330+331-305+351+338-309+310 Store in dark.  
 Concentrated Nitric Acid can decompose to nitrogen oxides (NO<sub>x</sub>) through action of heat or light resulting in a yellow colouration. However, this does not affect the performance of the acid with respect to trace metals or oxidising power. Storage in a cool, dark place is recommended.



### iso-Octane (see 2,2,4-Trimethylpentane)

iso-Pentanol (see iso-Amyl Alcohol)

Perchloroethylene (see Tetrachloroethylene)

Petroleum Distillate (see Petroleum Ether)

Petroleum Spirit (see Petroleum Ether)

iso-Propanol (see Propan-2-ol)

Propanone (see Acetone)

iso-Propyl Alcohol (see Propan-2-ol)

### Petroleum Ether 40-60°C tech

RS601

2½LT RS601L (Petroleum Distillate, Petroleum Spirit)  
 25LT RS601G BP 40-60°C d0.64 CAS[8032-32-4]  
 Dgr H:225-304-336-411-EUH066 Water <0.2%  
 P:210-233-243-273-280-301+310-303+361+353-304+340-331-403+235





















### Propan-2-ol tech

RS625

2½LT RS625L (iso-Propanol, iso-Propyl Alcohol)  
 25LT RS625G (CH<sub>3</sub>)<sub>2</sub>CHOH MW 60.10BP82.2°C d 0.78 CAS [67-63-0]  
 Dgr H:225-319-336 Assay >99% Water <0.5%  
 P:210-233-305+351+338



## ROMIL Technical Grade Solvents and Acids Specifications

<p>2½LT RA691L Dgr H:290-314 P:280c-301+330+331-305+351+338-309+310</p>	<p><b>Sulphuric Acid 1.84 tech</b> H<sub>2</sub>SO<sub>4</sub> MW98.07d1.84CAS[7664-93-9] Assay 95-98%</p>	<p>RA691 ☐</p>
	<p>Turpentine substitute (see White Spirit)</p>	
<p>2½LT RS702L 25LT RS702G Wng H:315-317-319-336-351-411 P:273-281-302+352-305+351+338-308+313</p>	<p><b>Tetrachloroethylene tech</b> (Perchloroethylene) CCl<sub>2</sub>CCl<sub>2</sub>MW165.83 BP 121.1°C d 1.62 CAS [127-18-4] Assay &gt;99% Water &lt;0.2% Unstabilised</p>	<p>RS702 ☐</p>
  		
<p>2½LT RS718L 25LT RS718G Dgr H:225-319-335-351-EUH019 P:210-240-305+351+338-308+313-403+233</p>	<p><b>Tetrahydrofuran tech</b> CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>OMW 72.11BP66.0°C d 0.89 CAS [109-99-9] Assay &gt;99%* Water &lt;0.3% *ex stabiliser Stabiliser: Butylated hydroxytoluene (BHT) ca. 250 ppm</p>	<p>RS718 ☐</p>
  		
<p>2½LT RS771L 25LT RS771G Dgr H:225-304-315-336-361d-373 P:210-240-301+310-331-302+352-403+235</p>	<p><b>Toluene tech</b> C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>MW92.14BP 110.6°C d 0.87 CAS [108-88-3] Assay &gt;99% Water &lt;0.2%</p>	<p>RS771 ☐</p>
  		
<p>2½LT RS901L 25LT RS901G Dgr H:225-304-315-336-410 P:210-233-240-273-301+310-331-302+352-304+340-403+235</p>	<p><b>2,2,4-Trimethylpentane tech</b> (iso-Octane) (CH<sub>3</sub>)<sub>3</sub>CCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub> MW 114.23 BP 99.2°C d 0.69 CAS [540-84-1] Assay &gt;98% Water &lt;0.2%</p>	<p>RS901 ☐</p>
   		
<p>2½LT RS864L 25LT RS864G Dgr H:226-302+312+332-304-315-319-334-411 P:261v-273-280f-301+310-331-305+351+338</p>	<p><b>White Spirit tech</b> (Turpentine substitute, Stoddard solvent) BP150-200°C d0.77 CAS[8052-41-3] Water &lt;0.2% Comprises a re ned mixture of mainly C9-C12 hydrocarbons being n-alkanes, iso-alkanes, cyclics and aromatics.</p>	<p>RS864 ☐</p>
   		
<p>2½LT RS982L 25LT RS982G Wng H:226-312+332-315 P:210-302+352-304+340</p>	<p><b>Xylene mixed isomers tech</b> C<sub>6</sub>H<sub>4</sub>(CH<sub>3</sub>)<sub>2</sub> MW106.17BP138-142°Cd0.86 CAS [1330-20-7] Water &lt;0.2% Comprises 3 isomers and ethylbenzene Assay (total C<sub>8</sub>H<sub>10</sub> isomers) &gt;97%</p>	<p>RS982 ☐</p>
